


AMERICAN ARACHNOLOGY

THE NEWSLETTER OF THE AMERICAN ARACHNOLOGICAL SOCIETY

No. 34

Nov. 1986



1987 MEETINGS

American Arachnological Society
to be held at Harvard University, Cambridge, Massachusetts

Dates: June 16, 17, 18 (Tues.-Thurs.) 1987. Field trip on June 19th.

Host: Dr. Herbert W. Levi, Invertebrate Department, Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts 02138

Housing: Graduate student dormitories, across Oxford Street from the Museum. Cost will be approximately \$27 per person per day. All rooms are singles. For many this is a steep fee; Those who have friends in the area may want to stay with them. On the other hand, commuting in and out of Cambridge can be a nuisance, whereas the dormitory is only a block from where the meetings will be held, and only a 5 minute walk from Harvard Square and the subway into Boston.

Parking: Daily parking permits for the Harvard University lots can be purchased for \$3 at the time of registration.

Registration: Tentatively set at about \$15 for students and \$25 for faculty. We are making every attempt to pare this down to offset the relatively high price of housing and eating out in Cambridge. The registration fee will cover the handbook and registration packet, lecture hall and any set up fees, security officer or a projectionist if needed, and the opening night dinner and social.

Food: People will be expected to pay for their own breakfasts, lunches, the mid-morning and mid-afternoon breaks, and Wednesday night dinner. The cafeteria lounge across from the lecture hall is reserved for our use, and the management is willing to stock up on our needs and preferences (i.e., cheap, good, interesting). For those who want to explore the many and varied eateries in Cambridge, the registration packet will include a map and suggestions, as well as indicating nearby supermarkets. In addition, the dormitory is equipped with kitchen and lounge facilities on each floor.

Anyone wanting to use these should bring their own cooking implements.

We anticipate that all will want to attend the Thursday night final banquet, a Boston Harbor cruise. Cost should not exceed \$20 for the 3-hour sunset cruise and buffet dinner.

Field Trip: Friday, June 19th to Beaver Brook Reservation, Hollis, New Hampshire.

Check-in and Check-out Times: People may arrive as early as Monday, June 15th; check-out is by 9:00 a.m. Saturday morning, June 20th. We realize that the meetings are brief; for many they also represent a blast from the past. Many participants may not have the time to visit old haunts and get in all the sightseeing they had hoped. We therefore encourage you, if you can afford it, to come a day early.

A pre-registration form is enclosed in the center of your newsletter. Please complete and send this form to: Invertebrate Department, Museum of Comparative Zoology, 26 Oxford Street, Cambridge, MA 02138

SOCIETY BUSINESS

Board of Directors Election: William Eberhard was elected to the Board of Directors in this spring's election.

Society Archives: At the annual business meeting the society voted to transfer its newly formed archives to the Smithsonian Institution. The Smithsonian will maintain existing materials and receive new items donated to the archives. Vincent Roth was reelected as Society Archivist. Anyone who has material to donate to the archives should send them to Vincent (Box G, Portal Arizona 85632) who will then organize and send it to the Smithsonian.

IN THIS ISSUE --

1987 MEETING - CAMBRIDGE, MA	1	Yael D. Lubin	3
SOCIETY BUSINESS ITEMS	1	David Wise	3
1988 MEETING	2	C. I. D. A. MEETING REPORT	3
BOOK REVIEW -- SPIDERS: WEBS, BEHAVIOR, AND EVOLUTION. W. A. SHEAR, EDITOR ...	2	A. A. S. 1986 MEETING: MEETING REPORT	3
NOTICES	2	FIELD TRIP REPORT	3
RESEARCH REPORTS:		STUDENT PAPER AWARDS	3
Joseph Beatty	2	INDEX TO ABSTRACTS	3
Norman Horner	3	ABSTRACTS	3

AMERICAN ARACHNOLOGY is the newsletter of the American Arachnological Society and is sent only to society members. For information on membership, write: Dr. Norman I. Platnick, Membership Secretary, American Arachnological Society, Department of Entomology, The American Museum of Natural History, New York, NY 10024, USA. Members of the Society also receive the JOURNAL OF ARACHNOLOGY.

Submission of items for AMERICAN ARACHNOLOGY or other correspondence concerning the newsletter should be directed to the editor, Dr. Brent D. Opell, Department of Biology, Virginia Tech, Blacksburg, Virginia, 24061, USA. Deadline for receipt of material for the fall issue of the newsletter is 24 September and for the spring issue, 24 March.

The society's executive committee recently voted in favor of the following items.

New Associate Editor of Journal of Arachnology: Jerome Rovner will replace William Peck as Associate Editor beginning 1 July 1987.

Anonymous Reviews Return: Effective immediately, reviewers for Journal of Arachnology are no longer required to sign their reviews. However, those who prefer to sign reviews may still do so.

Waiving Page Charges for Journal of Arachnology Articles: As announced earlier, the journal now has page charges for published articles. Decisions on reducing or waiving these charges for individuals unable to pay them will be made by the three Directors, using information supplied by the Associate Editor.

Exline-Frizzell Awards Committee: William Peck (Chairperson), Charles Dondale, and Matthew Greenstone will continue to serve on this committee for 1987. In 1988 the committee will consist of Gail Stratton (Chairperson), Charles Dondale, and Ann Rypstra.

1988 MEETING

The 1988 American Arachnological Society meeting will be held at New Mexico State University, Las Cruces, New Mexico. David Richman, Department of Entomology and Marsha Conley, Department of Biology will host the meeting, which will be held on 28, 29, and 30 June. Anyone interested in organizing a symposium should contact David Richman.

SPIDERS: WEBS, BEHAVIOR, AND EVOLUTION

Edited by WILLIAM A. SHEAR

This book, written by sixteen internationally recognized arachnologists, provides an up-to-date summary of what is known of the relationships of spiders to their webs, and the light cast by those relationships on the evolution of spiders--particularly as regards the constraints of behavior on evolution. The book confronts such fundamental questions as whether web-building spiders evolved from hunting spiders or vice versa.

There are contributions on web-site selection, webs as sensory transducers, and the effects of web geometry on prey-catching efficiency; and an entirely new field of research is opened by a chapter on how spiders take down their webs. Several chapters focus on orb-webs, those familiar and spectacular architectures of the garden, which provide insights into the evolutionary history of the spiders that weave them, and into what happens to spider behavior when spiders evolve "beyond the orb-web." Two chapters deal with the varied webs constructed by social spiders and how the web has facilitated and/or constrained the development of social behavior. Three others make detailed surveys of silk use in unusual groups of spiders (Mygalomorphae, Uloboridae, and Salticidae). The final chapter, a major new statement, summarizes the research on web evolution and stakes out new areas for future research.

There is an extensive taxonomic glossary to all families and genera of spiders mentioned in the text, with references to further information on their taxonomy and biology. The book is illustrated with 119 photographs (including many unusual web constructions) and 68 line drawings.

Shear, W. A. (editor). 1986. *Spiders: Webs, Behavior, and Behavior*. Stanford University Press, Stanford, California. 520 pages. ISBN 0-8047-1203-4. Price \$55.00.

CONTENTS

Introduction I

-- William A. Shear

1. Web-Site Selection: Are We Asking the Right Questions?

-- Anthony C. Janetos

2. Habitat Choice and Utilization in Web-Building Spiders.

-- Susan E. Riechert and Rosemary G. Gillespie

3. Transmission of Vibration in a Spider's Web.

-- W. Mitch Masters, Hubert S. Markl, and Anne J. M. Moffat

4. Effects of Orb-Web Geometry on Prey Interaction and Retention.

-- William G. Eberhard

5. Prey Specialization in the Araneidae.

-- Mark K. Stowe

6. Web Building and Prey Capture in the Uloboridae.

-- Yael D. Lubin

7. Social Spider Webs, with Special Reference to the Web of *Mallos gregalis*.

-- William James Tietjen

8. Web Building and Prey Capture in Communal Orb Weavers.

-- George W. Uetz

9. Web Building, Predatory Versatility, and the Evolution of the Salticidae.

-- Robert R. Jackson

10. The Role of Silk in Prey Capture by Nonaraneomorph Spiders.

Frederick A. Coyle

11. Web Removal Patterns in Orb-Weaving Spiders.

-- James Edwin Carico

12. The Monophyletic Origin of the Orb Web.

-- Jonathan Coddington

13. The Evolution of Web-Building Behavior in Spiders: A Third Generation of Hypotheses.

-- William A. Shear

Taxonomic Glossary, Bibliography, Index.

NOTICES AND REQUESTS

SPECIMENS NEEDED -- OPILIONES FROM CUBA

Luis F. de Armas (Academia de Ciencias de Cuba) and I are presently studying the Gagrellidae from the Greater Antilles and desire further materials and records from Cuba, Jamaica, and Hispaniola. We would also appreciate hearing from anyone (museum or individual) that has opiliones (all families) from Cuba for future studies. Please write to: James C. Cokendolpher, Department of Entomology, Texas Tech University, Lubbock, Texas 79409.

SICK ARACHNIDS??

Insect pathologist Carlo Ignoffo and I are trying to broaden our basic knowledge of *Nomuraea atypicola* and other arachnophilous fungi. If you find dead or dying animals with fuzzy or powdery coverings, we would appreciate receiving them for culturing. Please send them dry, (not in preservative!) to: Matt Greenstone, USDA, BCIRL, P.O. Box 7629, Columbia, MO 65205.

RESEARCH REPORTS

Joseph Beatty
Department of Zoology
Southern Illinois University
Carbondale, Illinois 62901

Right at the beginning of my work with spiders I quite accidentally became much involved with spiders on islands, and this interest continues.

My 25-year study of spiders on the Lake Erie islands has been terminated, and a paper on part of it is in press. I presented a discussion of the results at a Symposium on Biogeography at Ohio State University in summer of 1985.

During Dec. 1984 and Jan. 1985 I spent three weeks on Penang Island, Malaysia. With the help of a former student, Zairi Jaal, I was able to get about 60 *Liphistius*. I expected difficulty in locating their habitat or burrows, but there was none. They seem common on Penang. I found them every time I looked for them (not, of course, at every site I examined). Two brief trips to the mainland were less successful, though a few were found at Kedah Peak. Two specimens, an adult female of each of the two Penang species, are still living.

Last spring semester I taught a graduate level arachnology course for 3 students. One of them, Dan Mott, is working on a revision of North American mimetid, and another, Stephen Keffer, is doing non-thesis research on spiders with me. All four of us attended the Society meeting last summer.

At the end of December I will be going on sabbatical, and plan to join Jim Berry for more collecting in the Pacific. I also hope to get to Paris, London and Hamburg to examine museum material.

Norman Horner
Department of Biology
Midwestern State University
Wichita Falls, TX 76308-2099

It seems that the majority of my time is spent teaching and doing paper work; however, I do have a couple of "active" spider projects.

Cathy Tugmon, a masters' candidate, is doing her thesis research on spider karyotypes. She is primarily working with salticids and using 5 day old eggs. The technique followed is essentially that outlined by Matsumoto in 1977. (An observation of Somatic Chromosomes from Spider Embryo-Cells. *Acta Arachnol.* 27:167-172). To date, she has successfully karyotyped 8 species of salticids. If everything goes as planned, Cathy will be presenting the results next summer at the Harvard Meeting.

Dr. Fred Stangl, a colleague, and I observed a large population of *Salticus austiniensis* Gertsch under a rock ledge overhanging the shoreline of a local reservoir a couple of years ago. At that time, we were surprised to see so many in such close association. This past spring Fred discovered another population on the brick siding of his home. We certainly do not consider ourselves behaviorists; however, some interesting observations have been made. For example: 1) They appear to forage almost exclusively on vertical and inverted surfaces. 2) Intra-specific interactions are characterized by mutual avoidance and cannibalism appears to be nonexistent. 3) They feed almost exclusively on soft-bodied dipterans. 4) Where populations of *S. austiniensis* are found, *Metacryba undata* (DeGeer) appears to coexist with it. 5) Temperatures during our periods of observation ranged from 70°F to 102°F and seemed to have no major effect on spider activity. We anticipate that further observations will result in some useful and interesting notes on its biology.

We would be delighted to hear from anyone who could help us with these little guys.

Yael D. Lubin
Institute for Desert Research
Ben-Gurion University of the Negev
Sede Boqer Campus
84990, Israel

The move from Neotropical rainforest to Middle Eastern Desert involved many readjustments. On the 'plus' side, working in the canopy of a desert shrub habitat is certainly easier than in tropical rainforest canopy, and one needn't worry about stinging *Paraponera* ants, or ticks and mosquitoes. The 'minuses' are that field work is more strictly regulated by climate (only 'mad dogs and Englishmen go out in the midday sun'), and the lack of stinging and biting arthropods is compensated by the prevalence of highly poisonous snakes.

One cannot help but be amazed at the diversity of adaptations to this extreme desert environment. In general, the behavioral and life history adaptations of arthropods to deserts are poorly understood. The widow spiders, *Latrodectus*, seemed like a good place to start: There are many species in Israel, and some of these are strictly desert-adapted. (The systematics of the Israeli species is well known, thanks to a recent revision of the genus by G. Levy and P. Amitai). *Latrodectus revivensis* is one such species; it is apparently restricted to the Negev desert, but also overlaps with two other species, *L. pallidus* and *L. tridecimguttatus*, which have broader ecological and geographic distributions. My approach to studying the evolution of desert adaptations in this group is (1) to investigate the cause and effect relationships between environmental factors and behaviors, and (2) to compare behaviors among the three species.

L. revivensis and *L. pallidus* have similar web structures. These webs are specialized traps for ambulatory, terrestrial arthropods. However, the details of web structure, website selection and placement of webs differ in the two species, as do some aspects of their life histories. Since the web has many functions in the life of the spider, I contend that web design (structure, websites and placement) is critical and will be under strong environmental selection. Consequently, I have chosen to test ideas about the influence of environmental factors on behavior by examining web design (in the broad sense) in these two species, with the emphasis on mechanisms of adaptations in *L. revivensis*.

I am looking at adaptations relating to four main functions of the web: prey capture, microclimate control, reproduction, and protection from predators or parasites. (1) *L. revivensis* webs are located on isolated shrubs that may serve as foci for arthropod activity. I am investigating web and website characteristics, web orientation, and movement of spiders between websites with respect to food supply. (2) Webs of both species have cone or funnel-shaped retreats, placed high in shrubs, in which the spiders sit during the day. *L. revivensis* decorates the outer walls of the cone with prey remains, while *L. pallidus* uses stones, sand and twigs. Do these function in regulating the microclimate within the retreat? Preliminary measurements indicate that they do. (3) Males often inhabit webs of adult and subadult females, sometimes adding their small webs to those of the females. Movements of females between websites may be related to the presence or absence of males. In addition, "eggsac webs" of mature females are structurally different from webs of previous instars. The significance of these differences is not clear. (4) Protection from predators is perhaps the most difficult function to observe. Egg predators do exist. As with most of these functions, manipulations of webs and websites should yield answers to many of the questions.

David Wise
Department of Biological Sciences
University of Maryland
Baltimore County Campus
Catonsville, MD 21228

Currently, I am working on a book entitled "Spiders in Ecological Webs", to be published by Cambridge University Press as a contribution to the series Cambridge Studies in Ecology. In this book I will examine what we know about the position of spiders in ecological communities, simultaneously discussing and analyzing several broad issues in population and community ecology. One major focus, for example, will be the role of manipulative field experiments—their virtues and limitations—in ecological research. The book will be addressed to the general ecological community, the goal being to reach a diverse audience. In the process of treating topics of general ecological interest, I hope also to inform fellow ecologists about the importance and place of spiders in natural communities, perhaps winning converts to our ranks in the process. In order to make the book as current as possible when it is published, I would like to incorporate the most recent research on spider ecology. Thus I would appreciate receiving copies of any manuscripts you have in press and hearing about current research—your own, your students or that of colleagues.

Last month I returned from almost 15 months in Europe, having spent my sabbatical with Matthias Schaefer in Goettingen, West Germany. The year was highly productive and rewarding. I addition to collaboration and numerous discussions with Prof. Schaefer, I had the opportunity to visit and meet with many European arachnologists and ecologists. I initiated two projects (in addition to my book) while in Goettingen: 1) a field experiment to follow the decomposition rates of different types of leaf litter in a beech forest, and 2) development of a mathematical model of energy flow through the decomposer community that Prof. Schaefer, his students and colleagues have been studying for several years. I am re-orienting my research to the study of interactions in complex assemblages of species, and used my sabbatical to explore the feasibility of working with terrestrial decomposers and their natural enemies. I have not abandoned spiders, as they are major predators in such communities. My future research will deal with spiders as one compartment, albeit a major one, of the systems to which they belong.

Before going on sabbatical I initiated research on the intriguing color polymorphism exhibited by the theridiid *Enoplognatha ovata*. My Ph.D. student, Paul Reillo, is continuing this research with the support of a NSF Predoctoral Fellowship.

I hope to hear from many of you about ongoing ecological research that should be included in the book I am writing. I want to make it as complete and up to date as possible, and will appreciate any help you can give me in incorporating the latest findings.

1986 C.I.D.A. MEETING REPORT

by B. Opell

The tenth International Arachnological Congress, was held from 2-7 September 1986 in Jaca, Spain; a town of approximately 12,000, located in the Pyrenes of north central Spain. The meeting was hosted by Maria Rambla, Enrique Balcells, Jose Barrientos, Eugenio Ortíz, Cesar Pedrocchi, Carles Ribera, Carmen Urones, and Miguel Ferrandez.

The congress opened with a poster session, a reception hosted by Jaca's mayor, and a tour of the city. Each of the remaining four meeting days

began with a major address, included paper presentations, and concluded with a discussion. On the first day Rainer Foelix opened the meetings with a presentation on recent developments in spider neurobiology, on the second day Rainer Blanke discussed homologies in spider mating behavior and their usefulness in systematics, on the third day Patrick Blandin discussed the spatial and temporal structure of spider communities, and on the final day Jochen Martens summarized recent research in opilionid systematics, emphasizing the importance of genitalic characters. Concluding discussions focused on analogies and homologies in sensory and glandular structures of arachnids (lead by R. Legendre), ethospecies and ethological patterns as interspecific barriers (lead by L. Vlijm), spatial and temporal distribution patterns in arachnid fauna and methods for appraising them (lead by E. Duffy) and, methods of establishing serial relationships (lead by N. Platnick).

Jaca's many cafes and bars afforded opportunities for less formal discussions of these topics as well the 70 papers and 18 posters that were presented. Following the first two days of paper presentations, the congress excursion provided a chance to see surrounding regions. This excursion began with a visit to Santa Cruz de la Seros and a tour of its romanesque church. A winding road next took us to tenth century monastery at San Juan de la Pena and then to the high plane where a "new" monastery was build in the late seventeenth century to replace it.

The congress ended with an elaborate banquet, complete with an exhibition of traditional dancing. Many participants departed the following day, although about 30 stayed for a three-day, post-congress excursion that featured visits to villages and collecting stops in a variety of habitats. The next congress will be held in Turku, Finland and will be hosted by Pekka Lehtinen.

1986 A.A.S. MEETING

Meeting Report

by B. Opell

From 19 to 21 June, approximately 62 arachnologists gathered at Lindenwood College, St. Charles, Missouri for the society's 1986 meeting. Bill Tietjen, Alan Cady and L. Rao Ayyagari, all of Lindenwood, hosted the meeting and introduced its participants to the local night life. St. Charles is an old riverfront town on the Missouri River, not far from St. Louis. Lindenwood College entered the picture in 1827 and remains a pleasant, quiet campus even when invaded by arachnologists.

The 32 papers presented at the meetings gave participants a chance to learn about the latest findings of familiar research programs. For example, Allen Brady spoke about new problems and solutions in Lycosid systematics, Jonathan Coddington and Claudia Sobrevila told us that dinopids can use both forward and backward web strikes to capture prey, George Uetz informed us that there are really three species of colonial Metepeira, and Graeme Wilson showed that jumping spiders can see red. Many of the papers introduced new topics. Bret Beal showed that members of the fossil order Kustarachnida are really opilionids, Petra Sierwald demonstrated the usefulness of ontogenetic evidence in studies of spider genitalia, and Linda Rayor described the impact of damselfly predators on web site tenacity and reproductive success of pholcids.

Mat Greenstone raised our concern about spider hygiene by demonstrating their susceptibility to fungus and Jerry Rovner restored some peace of mind by showing that the setae and behaviors of many ground-dwelling spiders help prevent them from drowning during floods. As always, the creativity and quality of student papers provided good models for us all. For example, Jeff Schultz presented an elegant analysis of Dolomedes aquatic terrestrial locomotion, showing that leg coordination in spiders is achieved by ipsilateral rather than diagonal intersegmental nervous coupling. Giselle Mora described an intriguing opilionid species whose males construct nests and guard eggs.

The meeting concluded with a banquet and an evening session during which Lenny Vincent presented an illustrated tribute to B.J. Kaston. This session also included presentations of the latest para-arachnological research. Employing their well-known "deadness index", Bill Tietjen and Alan Cady discussing the ability of spiders to withstand submergence in various liquids and Jerry Rovner provided evidence for the monophyly of primates and arachnids.

Field Trip Report

by B. Vogel

Lincoln Hills, in the Cuivre River State Park, was the site of the field trip for the national AAS meeting. Lincoln Hills is a limestone highlands that drops eastwardly to the mighty Mississippi and has probably served as a refugium from the ice sheets. Biologically this area is very similar to the famous Ozarks located to the south and contains a great diversity of habitats: hardwood forests, eastern-type prairies, dry brushy ridges and deep cool ravines flowing with cool clear streams.

Our first stop was at Sac Prairie, one of the open grasslands managed by controlled burning. It was a gently undulating area of a few acres with an interesting diversity of vegetation. The spider fauna, alas, was not so diverse -- *Hycia*, *Metaphidippus* and *Dictyna*. It is my opinion that burning, which may be benign for small vertebrates, can be disastrous for arthropods.

The next brief stop was at one of the cool ravines with a clear flowing stream (Little Sugar Creek?). Collecting here was also on the meager side, as the creek had recently been scoured by a flood which rose to 6-10' above stream's current level. Those who went upstream said they found *Pardosa*, but I went downstream and found nothing on the gravel bars, or under rocks. I found *Xysticus* and *Micrathena* high up on the greenery, until I was blocked by the dread P.I.

The final stop at a picnic shelter gave access to a hardwood forest, forest creek, open sunny forests and dry brushy ridge trail. We all scattered in different directions, collecting merrily until lunch time. The catch included *Neoantistia*, *Dolomedes*, *Lecaue*, gnaphosids, *Microthema*, *Teutalena elegans*, *Oxyopes scularis*, *Phidippus johnsoni*, *Tetragnatha*, *Mimetus*, *Philodromus*, *Argyrodes*, *Steatoda*, *Pirata*, *Tegenaria* and opilionids. Not a bad sample for a half day outing. I would have preferred a few more hours in Cuivre River Stat Park, but the bus had to return to the barn.

STUDENT PAPER AWARDS

This year's first place student paper award was given to Jeffry W. Schultz, Department of Anatomy, University of Chicago for his paper entitled "Walking and surface film locomotion in terrestrial and semi-aquatic spiders." Giselle Mora, Department of Zoology, University of Florida, received the second place student paper award for her paper entitled "Nest Construction and nest acquisition strategies of daddy long legs with male egg guarding behavior."

INDEX TO ABSTRACTS

- Ecology & Population Biology: 8, 10, 11, 14, 16, 17, 19.
Ethology & Neurobiology: 3, 5, 8, 11, 12, 18, 19, 21, 24, 25, 26.
Morphology & Physiology: 4, 6, 7, 9, 13, 15, 18, 22.
Evolution & Systematics: 1, 2, 12, 15, 20, 23.
History of Arachnology: 28.

ABSTRACTS

Bret S. Beal (University of Michigan) 1
REINTERPRETATION OF THE KUSTARACHNIDA.

Petrunkévitch erected the order Kustarachnida to accommodate three poorly preserved Carboniferous arachnid fossils. Re-examination of these specimens indicates that several features cited as diagnostic of kustarachnids (pedicel; chelate pedipalps; pedipalpal coxae fused medially; double trochanters) are misinterpretations of the fossil material. Other characters (e.g., triangular coxae; legs long and slender; two eyes on median tubercle) suggest that these fossils are actually opilionids. One feature, a three-segmented "pygidium" on one specimen, is not observed in extant opilionids, but it may be plesiomorphic for the order. The name Kustarachnida should be regarded as a nomen nudum. This reinterpretation is pertinent to clarifying arachnid ordinal relationships.

Allen R. Brady (Hope College) 2
SYSTEMATIC PROBLEMS IN THE LYCOSINAE

Over 50 species of medium to large size wolf spiders from the Nearctic Region have been placed in the genus Lycosa. Recent studies indicate that several distinct genera are included in Lycosa. Roewer (1958, 1959) established 51 new genera of Lycosinae based primarily upon differences in

the number of posterior cheliceral teeth and the arrangement of the eyes in the anterior row. My investigations of North American Lycosinae indicate that color patterns on the dorsal surface of the carapace, length of legs relative to body size, and the structure of the male and female genitalia are most reliable in determining generic relationships. The new lycosid genus *Gladicosa* has been established based upon these characteristics. Measurements of eye arrangement, leg lengths, and bodily dimensions are not necessary for differentiating species, but they are very useful in establishing species groups and genera.

John A. Bruce (Washington State Univ.) & James C. Carico (Lynchburg College) 3
SEXUAL BONDAGE IN SPIDERS: THE MATING BEHAVIOR OF PISAURINA MIRA (WALICK.) (PISAURIDAE)

The mating behavior of the nursery web spider *Pisaurina mira*, is described for the first time. These spiders mate while suspended from draglines as in *Oxyopes heterophthalmus* (Latr.) and *Peucetia viridens* (Hentz). The unique feature of the mating in *P. mira*, however, is the male's use of a veil of silk to wrap the female's legs I and II into a folded position. Mating is accomplished in a version of position II with bodies in a right angle as the female is cradled by the male's legs. The use of silk to "tie" the female is known elsewhere only in *Xysticus*.

James E. Carrel (University of Missouri-Columbia) 4
THE SPIDER HEART AS A PHYSIOLOGICAL INDICATOR

Evidence that a spider's heart is a major indicator of its physiological condition will be reviewed. The influence of exercise, temperature, dietary toxins, and other stimuli on spider cardiac frequencies will be presented. Finally, the controversy over the relationship between resting heart rates and resting metabolic rates in spiders will be discussed.

Jonathan Coddington and Claudia Sobrevilla (Smithsonian Institution) 5
TWO STEROTYPED ATTACK BEHAVIORS IN DINOPIS SPINOSUS MARX.

A series of experimental manipulations documents that *Dinopis spinosus* Marx has at least two stereotyped attack behaviors. In "backward" strikes the spider expands the net backward over its cephalothorax and away from the substrate; these strikes capture aerial prey. Vibratory stimuli are necessary and sufficient to elicit backward strikes. In "forward" strikes the spider expands the web downward and forward against the substrate; these strikes capture walking prey, and visual rather than vibratory stimuli are more effective. The mechanisms enabling each type of strike and the extraordinary extensibility of the web are described. This work resolved a long-standing controversy in the literature over the manner in which dinopids catch prey.

Matthew H. Greenstone, Carlo M. Ignoffo (U.S.D.A., Biological Control of Insects Research Laboratory, Columbia, Missouri) and Robert A. Samson (Centraal Bureau Voor Schimmelcultuur, Baarn, The Netherlands). 6
SUSCEPTIBILITY OF SPIDER SPECIES TO THE FUNGUS, NOMURAEA ATYPICOLA.

The fungus *Nomuraea atypicola* has been associated with spiders but reported only sporadically from a few species in anecdotal field accounts. We tested twenty species in ten families of spiders and one species of harvestman for susceptibility to the fungus. Our results show that spiders, and possibly arachnids generally, are widely susceptible to the fungus.

Craig S. Hieber (University of Florida) 7
MORE CONTAINER SCIENCE: WATER, FUNGUS, AND DROWNING

The ability of the cocoons of *Mecynogea lemniscata* and *Argiope aurantia* (Araneidae) to control fungal attack and drowning were examined. The results of laboratory and field experiments indicate that the component

parts of these cocoons do not conduct water toward or into the cocoon. How this relates to fungal attack is unclear. The cover and suspension system of *M. lemniscata* cocoons play no role in controlling fungal attack in the egg stage, although the cover is important in the spiderling stage. The suspension system of this cocoon is also important in protecting eggs and spiderlings from drowning. In contrast, the cover and suspension system of *A. aurantia* cocoons play no role in protecting either the eggs or spiderlings from fungal attack or drowning. The observed differences between the two cocoons may be related to cocoon size and the thickness of the flocculent silk layer within the cocoons.

Maggie Hodge (University of Cincinnati) 8
THE RELATIONSHIP BETWEEN AGNOSTIC BEHAVIOR AND SPATIAL ORGANIZATION IN DESERT AND TROPICAL METEPEIRA SPP. FROM MEXICO.

Populations of *Metepeira* spp. from desert and tropical regions of Mexico show varying degrees of social behavior. Recent work suggests that differences in spacing behavior are genetically controlled and that spacing seems to be maintained by aggressive interactions between individual spiders. This research examined the relationship between agonistic behavior and spacing in populations of these spiders raised under identical conditions of space and food availability in the laboratory. These studies found that individuals from the desert population show a greater tendency to behave agonistically towards conspecifics, whereas the tropical spiders show a high level of tolerance for one another. Interaction strategies used in disputes over web sites were found to differ in tendencies to escalate to conflict and to abandon a fight. Continued studies in the field should give us some ideas about how different behavioral strategies can evolve in different environments.

J. K. Johnson (University of Kansas) 9
SHEDDING OF PHOTORECEPTOR MEMBRANE IS DEPENDENT UPON LIGHT IN THE OPILIONID RETINA.

Shedding of photoreceptor membrane occurs over a 24-hour period in *Mesosoma roeweri* (Opilionida: Palpatores) and *Vonones ornata* (Opilionida: Laniatores). Animals from both species were dark-adapted (DA) for periods of 2, 26, 50, 74, and 98 hours beyond normal light onset. Additional animals were DA 98 hours then given 30 minutes of light; 30 minutes of light followed by 60 minutes of dark, or 90 minutes of light and fixed. Results indicate two spatially distinct shedding pathways exist in *V. ornata* (intracellular and intercellular), whereas only one pathway exists in *M. roeweri* (intracellular). The intracellular pathway is attenuated by the prolonged absence of light in both species. However, intercellular shedding in *V. ornata* is unaffected by the prolonged absence of light, but is reset to zero and the rhabdom is structurally rejuvenated by a 30 minute pulse of light.

David A. Landes, James Hunt (University of Missouri-St. Louis), and Alan B. Cady (Lindenwood College) 10
SEASONAL AND LATITUDINAL VARIATION IN SPIDER PREY OF THE BLUE MUD DAUBER CHALYBION CALIFORNICUM

Spider prey of the Blue Mud-Dauber, *Chalybion californicum* (Saussure) in eastern Missouri included 15 species representing 4 families. Araneidae was the most diverse family, with 9 species, but Theridiidae was more numerous, with *Steatoda americana* (Emerton) outnumbering all other species combined. Theridiidae constitute a majority of the total individuals in both this survey and a similar survey from Maryland; corresponding studies in southern Oklahoma and Florida yielded high proportions of Araneidae. The present study resembled the two southerly sites in having more species of Araneidae than any other family, while the most northerly survey found more species of Theridiidae. Latitudinal patterning of these differences suggests that analysis of the spider prey of mud-dauber wasps may provide insights into the biogeography of spider diversity.

Gary L. Miller and Patricia Ramey Miller (Weber State College) 11
REPRODUCTIVE BIOLOGY AND COURTSHIP BEHAVIOR OF GEOLYCOSA TURRICOLA (TREAT).

The results of field and laboratory studies of the reproductive cycle and courtship behavior of the obligate burrowing spider *Geolycosa turricola* are reported, and comparisons are made to the reproductive patterns of other lycosids. Some components of the typical lycosid male courtship, such as palpal displays and palpal drumming, are absent or reduced. Modifications in other displays are described. A cohabitation phase (immature female with mature male) is described and its significance discussed. *Geolycosa* reproductive biology is discussed in terms of limitations related to burrowing.

Giselle Mora (University of Florida) 12
NEST CONSTRUCTION AND NEST ACQUISITION STRATEGIES OF DADDY LONG LEGS WITH MALE EGG GUARDING BEHAVIOR.

Zygopachylus albomarginis (Opiliones:Gonyleptidae) is the only arachnid known to have paternal care. Males build nests out of mud and tree bark or take over previously-constructed nests. Females court males and lay eggs inside the nests. I will present observations on the natural history and reproductive biology of this species and will discuss nest acquisition strategies of males under different ecological conditions.

Brent D. Opell (Virginia Tech) 13
CHANGES IN FORCE ASSOCIATED WITH WEB REDUCTION IN THE SPIDER FAMILY ULOBORIDAE.

Resting and maximum force measurements were taken of a developmental series of *Hyptiotes cavatus*, *Uloborus glomosus*, *Miagrammopes animotus*, *Miagrammopes pinopus*, and an undescribed Costa Rican *Miagrammopes* as they hung from a thread spun between the glass needle and fixed support of a strain gauge. Both carapace length and spider weight were used as an index of size. When regression analyses of resting and maximum forces were compared, *H. cavatus* exerted the greatest relative force and *Miagrammopes* species the least. Within the genus *Miagrammopes*, the species with the most highly modified carapace expressed the greatest force.

William J. Pfeiffer (Center for Energy and Environmental Research, University of Puerto Rico) 14
COMMUNITY STRUCTURE AND POPULATION DYNAMICS FOR SPIDERS OCCUPYING THE LITTER LAYER OF A PUERTO RICAN RAINFOREST

Litter arthropods from a subtropical wet forest in the Caribbean National Forest were censused with a D-Vac suction sampler at roughly 40-day intervals for one year. The 40 samples collected at each interval were subjected to Tullgren extraction and manual sorting. The spider community was characterized by: a) a low species richness (20 species from 12 families), b) high community densities (mean annual density - $354 m^{-2}$), c) numerical domination by two species, a pholcid and an ochyroceratid, which account for 62% of over 4300 specimens, d) the lowest annual variability of community density in available literature, and e) reduced body size for the community as a whole, suggesting high turnover rates and production/biomass ratios. Spider densities were significantly related to litter standing crop and potential prey densities. The high proportional representation of spiders in this arthropod community has also been noted for other Puerto Rican habitats (secondary growth vegetation) in previous studies. Standing stock estimates based on morphometric regressions and preliminary production values from the size-frequency method will be presented.

Norman I. Platnick (American Museum of Natural History) 15
THE TIBIAL AND PATELLAR GLAND OF THE LETTONETIDAE

Linda S. Rayor (Department of Systematics & Ecology, University of Kansas, Lawrence, Kansas) 16
EFFECTS OF DAMSELFLY PREDATION ON PHOLCID SPIDER FITNESS AND WEB SITE TENACITY.

The large neotropical damselfly, *Mecistogaster modestus*, preys exclusively on small web-building spiders in the rainforest. One pholcid spider species, *Modisimus* sp. C, receives 55.3% of the damselfly predation attempts, but escapes from 51% of those attacks. Pholcids in sunflecks and gaps are subject to much greater predation risk than pholcids living in shadier areas with a denser canopy, because the damselflies only forage in direct sunlight. However, the insect prey available to the spiders appears to be greater in the sunnier areas. I am examining the effects of the conflicting pressures of predation risk and food acquisition on the pholcids' web-site tenacity and reproductive fitness. I will discuss aspects of this predator-prey system, and results from my first field season.

Jon Reiskind (University of Florida) 17
LYCOSA ERICETICOLA, A NATIVE FLORIDAN WOLF SPIDER

The entire range of the Rosemary Wolf Spider, *Lycosa ericeticola* Wallace, is a compact area of about 2500 hectares in north central Florida. Its distribution is almost entirely found within that of the evergreen shrub Rosemary, *Ceratiola ericoides*. The distinctiveness of the male genitalia of *Lycosa ericeticola* makes it easy to distinguish from its congener and close relative, *L. ammophila*. The overall appearance of these two species is very similar (if not identical) and both are found in the Rosemary habitat but they have never been found sympatric. The cause of this exclusionary pattern of distribution is not known but may be historical or due to some sort of competitive interaction. A third member of the species group, *L. timuqua*, is found in association with both of these species although with a slight difference in preferred substrate (on leaves rather than open sand).

Jerome S. Rovner (Ohio University) 18
DENSE SETAL DISTRIBUTION AND REDUCED LOCOMOTION ENHANCE FLOODING SURVIVAL IN INLAND TERRESTRIAL SPIDERS

Three of six species of spiders collected from webs under stones showed no apparent effects of a 1-hr submergence. These spiders had dense coats of hydrofuge hairs on the opisthosoma that trapped an extensive air store. This bubble prevented entry of water into the respiratory openings and provided a small oxygen supply and short-term physical gill. The hydrofuge hairs also enabled the spiders to float high and dry on the surface after being buoyed up by the bubble. The slower and reduced locomotion of these spiders during submergence probably further reduced their already low oxygen needs. Four aerial web weavers fared poorly when submerged, especially two araneid species, which had sparse setae yielding little or no air store. These and previous data suggest that setal density in terrestrial spiders relates partly to drowning resistance.

Ann L. Rypstra (Miami University, Hamilton Ohio) 19
A COMPARISON OF THE FORAGING SUCCESS IN SOLITARY AND AGGREGATED SPIDER WEBS

Recent foraging theories have taken into account both the mean and variance associated with food intake. These theories predict that hungry individuals should select the foraging mode with the greatest variance (risk prone) so that they have some probability of meeting their metabolic needs. Observation of foraging success in spiders indicate that mean capture success is similar between individuals and aggregates. However, variability in daily food intake is much greater for solitary spiders than for those in groups. Similar results were obtained in prey introduction experiments. Insects entering groups were frequently captured by the second or third spider they encountered, whereas spiders in lone webs either captured their prey or lost it. Since spiders in the temperate zone only seem to aggregate in areas of high prey, these results are consistent with risk sensitive foraging theories.

William A. Shear (Hampden-Sydney College) 20
EVOLUTION OF THE OPILIONID SUPERFAMILY ISCHYROPSALIDOIDEA

A cladistic analysis of the opilionid superfamily Ischyropsalidoidea revealed three groups of genera, each probably monophyletic, and recognized here as families. The family Ischyropsalididae Simon is redefined to include only the genus *Ischyropsalis*, the family Sabaconidae Dresco is expanded to include *Sabacon* (= *Tomicornerus*, new synonymy) and *Taracus*, and the family Ceratolasmatidae is described as new for the genera *Ceratolasma*, *Hesperonemstoma*, *Crosbycus*, and *Acuclavella*, new genus.