

ISSR (Inter Simple Sequence Repeats) as molecular markers to study genetic diversity in tarantulas (Araneae, Mygalomorphae)

Salima Machkour-M'Rabet^{1,4}, Yann Hénaut¹, Ariane Dor¹, Gabriela Pérez-Lachaud², Céline Péliissier³, Charles Gers³ and Luc Legal^{3,4}: ¹Ecología y Conservación de Fauna Silvestre, El Colegio de la Frontera Sur, Avenida Centenario Km 5.5, AP 424, 77900 Chetumal, Quintana Roo, Mexico; ²Entomología Tropical, El Colegio de la Frontera Sur, Carretera Antigua Aeropuerto Km 2.5, Apdo. Postal 36, 30700 Tapachula, Chiapas, Mexico; ³ECOLAB (Laboratoire d'Ecologie Fonctionnelle), UMR 5245 (CNRS-UPS-INPT), Université Paul Sabatier, Bâtiment IRV3, 118 Route de Narbonne, 31062 Toulouse cedex 4, France; ⁴Departamento de Sistemática y Evolución, CEAMISH-Universidad Autónoma Estado Morelos, Cuernavaca, Morelos CP: 62210 México

Abstract. Although all species of the *Brachypelma* genus are protected under CITES, few studies have been performed on the genetic structure of the populations of these endangered tarantulas. Here we propose, for the first time in spiders, to use ISSR (Inter Simple Sequence Repeat) technique to study the genetic variability of Mexican populations of *Brachypelma vagans* (Ausserer 1875). We used a nonlethal technique to collect samples from six populations in the Yucatan peninsula and we tested seven ISSR primers. Four of these primers gave fragments (bands) that were sufficiently clear and reproducible to construct a binary matrix and determine genetic variability parameters. We revealed a very high percentage of polymorphism ($P = 98.7\%$) the highest yet reported for tarantula spiders. Our results show that the ISSR-PCR method is promising for intraspecific variation of tarantula spiders.

Keywords: ISSR, Theraphosinae, *Brachypelma*, genetic population, Mexican redrump tarantula

Members of the genus *Brachypelma* are charismatic spiders, being colorful, large, and docile (Locht et al. 1999). The pet trade, habitat destruction, high mortality rates as juveniles, and late sexual maturity result in all *Brachypelma* species being listed in Appendix II of CITES. In recent years, efforts have been made to increase knowledge of their ecology (Yáñez & Floater 2000; Machkour-M'Rabet et al. 2005, 2007) and behavior (Locht et al. 1999; Reichling 2000). However, studies to better understand the genetic structure of tarantula populations are essential to assess the conservation status of the genus. Recently, the development of molecular techniques has helped inform conservation strategies.

Here, we focused our effort on *Brachypelma vagans* (Ausserer 1875), which is distributed from Southern Mexico south to Costa Rica (Locht et al. 1999), but has also been recorded outside its natural range in Florida as a result of the release of pet trade animals (Edwards & Hibbard 1999). As with the study of most tarantulas, the biology and ecology of *B. vagans* is poorly known (Carter 1997; Yáñez et al. 1999; Machkour-M'Rabet et al. 2005, 2007) and little information exists on the genetic structure of its populations (Longhorn et al. 2007).

Mitochondrial DNA and allozyme electrophoresis have been used previously to evaluate population genetics in Mygalomorphae (Ramirez & Froehlig 1997; Bond et al. 2001; Pedersen & Loeschcke 2001; Ramirez & Chi 2004; Bond et al. 2006; Arnedo & Ferrández 2007). For the *Brachypelma* genus (Theraphosidae) only one study has been carried out recently (Longhorn et al. 2007), which focused on the genetic structure of two Belizean populations of *B. vagans* using two portions of mitochondrial DNA (partial 16SRNA + tRNA-Leu + partial ND1 and CO1) and one nuclear non-coding gene (ITS-2). This study showed that nuclear markers are relatively invariant across *B. vagans* populations while mitochondrial

markers possess sufficient resolution to estimate the genetic structure of this species. However, it has been suggested that alternative sources of nuclear genes could be used to enhance the characterization of population structure in tarantula spiders. In this context, and because no microsatellite primers are available for *Brachypelma*, we here explore the usefulness of a relatively novel technique in animals, Inter Simple Sequence Repeats (ISSR), to discriminate among populations.

Dominant ISSR markers are widely used in the conservation of rare plants (Kothera et al. 2007) and are being increasingly used in animals (Wink et al. 2002; Hoffman et al. 2006; Guicking et al. 2006; Joger et al. 2007), particularly invertebrates (Abbot 2001; Luque et al. 2002; Chatterjee & Mohandas 2003; Hundsdoerfer & Wink 2006; Roux et al. 2007). However, until now, this technique has not been applied to spiders.

The PCR-ISSR method was used here to screen a large part of the genome without prior knowledge of the sequences. This provides highly reproducible results and generates abundant polymorphisms. The great advantage of ISSR is that the primers work universally for many animal and plant species. Consequently, it is not necessary to define PCR primers for each species, unlike microsatellites. Furthermore, ISSR demands fewer experimental steps and is therefore easy to carry out with a low cost-benefit ratio compared with RFLP (Restriction Fragment Length Polymorphism) and results in a higher reliability and repeatability than RAPD (Random Amplification of Polymorphic DNA; Nagaraju et al. 2001; Luque et al. 2002). Absence of a band is interpreted as the loss of a locus through either the deletion of the SSR (Simple Sequence Repeat) site or a chromosomal rearrangement (Wolfe & Liston 1998). ISSR are thus considered and treated as dominant markers (Casu et al. 2005).

The method uses polymerase chain reaction (PCR) with repeat-anchored or non-anchored primers to amplify DNA sequences between two inverted SSR (Zietkiewicz et al. 1994).

⁴ Corresponding author. E-mail: smachkou@ecosur.mx




Figure 1.—Location of the six samples used in the study in the Yucatan Peninsula (represented by stars). The geographic distribution of the Mexican Redrump Tarantula, *Brachypelma vagans*, in Mexico is represented on the smaller map (upper left).

Therefore, the only DNA stretches amplified are positioned between two identical but inverted microsatellites (SSR). A single primer can amplify up to 80 loci simultaneously. This method provides genomic information available for a broad spectrum of applications: population genetics, hybridizations, and gene mapping (Wink 2006). The ISSR method represents one of the most promising tools in population genetics studies and deserves increased attention (Behura 2006).

The aim of this study was to technically adapt ISSR-PCR for tarantulas and to provide a preliminary assessment of whether this method is advantageous to explore the genetic structure of tarantula populations. The only previous study using dominant markers (RAPD; Hettle et al. 1997) to study genetic populations of tarantula spiders (*Brachypelma albopilosum* Valerio 1980) showed that none of the six primers used were reliable to differentiate inter- and intra-family relationships. Here we report the technical aspects of a recent molecular tool to study populations of endangered tarantulas.

METHODS

Brachypelma vagans samples were collected during March and April 2007 around six traditional villages of the Yucatan Peninsula (Mexico): Ley de Fomento: 18°03'N, 89°25'W; Conhuas: 18°32'N, 89°55'W; Zoh-Laguna: 18°35'N, 89°24'W;

Luis Echeverria: 18°39'N, 88°13'W; Raudales: 18°42'N, 88°15'W; and Cozumel: 20°21'N, 86°59'W (Fig. 1). We collected the samples using a nonlethal technique that consists of inducing limb autotomy (Longhorn 2002). In response to pressure, the limb will detach and the muscles will contract to prevent hemolymph loss; spiders in which a limb is removed will regenerate the limb during subsequent molts. We chose to remove the medial limb (III) because the anterior legs (I and II) are used in sensory behaviors and the posterior legs (IV) are used defensively in brushing urticating hairs (Smith 1994). Samples were preserved in 95% ethanol at room temperature, and sent for DNA analysis under CITES export permit (MX34176) to ECOLAB (Laboratoire d'Ecologie Fonctionnelle) at University Paul Sabatier (Toulouse, France).

A small part of the limb was cut off and incubated for 12 h at 50°C in 350 µl of buffer B (10 mM Tris, pH 7.5, 25 mM EDTA, and 75 mM NaCl) with 500 µg of proteinase K and 20 µl of SDS (20%). Proteins and residues were precipitated with 200 µl of saturated NaCl solution and centrifuged at 14,000 rpm for 30 min. DNA from the supernatant was saved and precipitated with 400 µl of cold isopropanol, mixed and centrifuged at 14,000 rpm for 40 min at 2° C. The isopropanol was eliminated and the precipitate was washed with 500 µl of 70% ethanol and centrifuged at 14,000 rpm for 10 min at 2° C. The precipitate was dried and redissolved in 100 µl of TE buffer (pH = 7) and preserved at -28° C until utilization. The concentration of the DNA obtained was determined by spectrophotometry (NanoDrop ND-1000) and the quality was checked using electrophoresis in agarose/TBE (1.2%) gel.

Inter Simple Sequence Repeat (ISSR) analysis was performed using seven primers (Table 1). PCR amplifications were performed in a 25 µl reaction volume containing ~20 ng of template DNA, 50 µM of primer (Invitrogen), 0.2 mM of each dNTP from dNTP Mix (Promega), 2.5 µl of 5× Green Buffer (Promega), 3 µl of MgCl₂ (1.5 mM, Promega), and 2.5 U of *Taq* polymerase (Promega). All amplifications were done in a T3 Thermocycler (Biometra). The cycling conditions were as follows: initial denaturation step at 94° C for 4 min, 39 cycles of denaturation at 94° C for 45 s, primer annealing at 56° C for 45 s, and extension at 72° C for 2 min, followed by a final extension at 72° C for 10 min.

Electrophoresis was performed with 7 µl of amplified products on a 2% agarose gel using 1× Tris acetate EDTA buffer at 140 V for ~2 h. The bands were detected with ethidium bromide under UV light and digitized (Bio-Vision 3000, Vilbert-Lourmat) (Figure 2).

In our first experiments, conditions for ISSR with different primers were not optimal. One of the most important factors is

Table 1.—SSR primers screened for ISSR-PCR in the tarantula *Brachypelma vagans*. B = T, C or G; D = A, T or G; R = A or G; Y = C or T; W = A or T.

Code	Sequence (5'→3')	Abbreviation	Amplification pattern	Total bands
CA	CACACACACACA	(CA) ₇	Poor amplification	-
CA+	CACACACACACARY	(CA) ₇ RY	Smear	-
+CA	RYCACACACACACA	RY(CA) ₇	Smear with band	-
ACA+	ACAACAACAACABDB	(ACA) ₅ BDB	Good	16
+ACA	BDBACAACAACAACA	BDB(ACA) ₅	Good	25
GACA+	GACAGACAGACAWB	(GACA) ₄ WB	Good	15
+GACA	WBGACAGACAGACACA	WB(GACA) ₄	Good	20




Figure 2.—Example of polymorphic ISSR banding patterns with one marker (+ACA) for two different populations: Cozumel (A) and Luis Echeverrerria (B).

the difference in the amount of DNA loaded that can weaken the quality of the electrophoretic resolution. In order to obtain comparable and reliable results, we used the same DNA concentration for all samples. We found a minimum optimal amount of 20 ng per sample. Another important parameter is the primer annealing temperature. We experimented with temperatures ranging from 46° C to 66° C with a step of 1° C for all primers and we chose an optimal temperature (56° C) identical for all primers to facilitate the PCR procedure. Also, we checked various parameters to optimize our results. The standard number of reamplifications (39 cycles) was used and gave repeatable and reliable results for all primers. The concentration of MgCl₂ was tested from 2.6 µl to 3.4 µl in steps of 0.2 µl and we found good results (quality of the electrophoretic resolution) for values above 3 µl. The concentration of buffer was checked from 2.1 µl to 2.9 µl in steps of 0.2 µl and these modifications had no influence on the results, then we chose a medium value of 2.5 µl. Finally, the method used for storage of the spiders limbs [i.e. preservation in ethanol (95%) and dry-conservation at room temperature (3 years old)], was checked. Only preservation in ethanol resulted in amplification.

Table 2.—Genetic diversity of *Brachypelma vagans* in the Yucatan Peninsula based on ISSR markers. *n*: number of individuals kept for analysis; N¹: number of bands scored; N²: number of polymorphic bands; N³: number of signature bands; *P*: percentage of polymorphism; *h*: Nei's gene diversity; *H*: Shannon Index; SD: standard deviation.

Population name	<i>n</i>	N ¹	N ²	N ³	<i>P</i> (%)	<i>h</i> (± SD)	<i>H</i> (± SD)
Raudales	22	64	61	1	80.26	0.273 (0.181)	0.411 (0.252)
Zoh-Laguna	24	65	58	0	76.32	0.272 (0.191)	0.405 (0.268)
Ley de Fomento	26	69	64	2	84.21	0.296 (0.178)	0.442 (0.243)
Conhuas	23	65	59	3	77.63	0.271 (0.196)	0.403 (0.270)
Luis Echeverrerria	26	64	56	1	73.68	0.284 (0.197)	0.418 (0.277)
Cozumel	27	56	43	0	56.58	0.193 (0.203)	0.288 (0.291)

The gel separation of ISSR fragments (bands) was used for each individual and each primer to score the presence (1) or absence (0) of bands. This information generated the binary matrix used for analysis. Only bands that could be scored consistently among populations were used, and we assumed that each marker band represented a distinct locus.

The binary matrix was used under Hardy-Weinberg equilibrium to determine the genetic diversity: percentage of polymorphism (*P*), Nei's gene diversity (*h*) using corrected allele frequency (Lynch & Milligan 1994) and the Shannon Index (*H*) (Lewontin 1972), at the species level and for each population. All analyses were carried out using POPGEN Version 1.32 (Yeh et al. 1997). In order to describe the genetic structure and variability among and between populations, non-parametric Analysis of Molecular Variance (AMOVA) (Excoffier et al. 1992) was performed with GENALEX V6 (9999 permutations; Peakall & Smouse 2006).

RESULTS

Of the seven primers initially tested for the six populations, only four produced clear reproducible fragments (Table 2). Interestingly, the most classic and polymorphic primer (CA_n)

for butterflies (Nagaraju et al. 2001; Luque et al. 2002; Hundsdoerfer & Wink 2006; Roux et al. 2007) failed in the tarantula. From these four primers, a total of 76 scorable ISSR fragments were selected in the 180 individuals screened from all populations (30 individuals for each population). In Table 2, the number of bands and the number of polymorphic bands for each population is given. In addition, we give the number of bands found only within each of the populations, which we call “diagnostic bands” (Table 2; Luque et al. 2002). The very low number of these bands indicates that all populations belong to the same species.

Of the 30 individuals of each population, we only kept individuals presenting a banding pattern for the four primers and for which the interpretation of the banding pattern was unequivocal. For this reason, the number of individuals used for analysis is lower than the number screened (Table 2).

The percentage of polymorphic loci (P) varied between populations (Table 2), ranging from 57% in the Cozumel island population to 84% in the Ley de Fomento population. A mean P of 98.7% was observed across the 6 populations. Nei's gene diversity (h) was low in the Cozumel population with 0.193 (SD = 0.203), while it was higher but relatively constant for continental populations (Table 2). The mean for all populations was 0.324 (SD = 0.164). For the Shannon Index (H), we observed a similar pattern (Table 2): Cozumel island diversity was lower, with a mean across all populations of 0.485 (SD = 0.213).

AMOVA analysis revealed that 79% ($df = 142$, $P < 0.001$) of the variability occurred among individuals within populations and that a strong genetic difference among populations was observed (21%, $df = 5$).

DISCUSSION

Our study revealed a high level of polymorphism for tarantulas in comparison with other studies using allozymes [$P < 7.7\%$ for *Aptostichus simus* Chamberlin 1917 (Cyrtauchenidae), Ramirez & Froehlig 1997; $P < 33\%$ for *Atypus affinis* Eichwald 1830 (Atypidae), Pedersen & Loeschcke 2001; $P < 30\%$ for *Antrodiaetus riversi* (O. Pickard-Cambridge 1883) (Antrodiaetidae), Ramirez & Chi 2004 (reported therein as *Atypoides riversi*)]. However, the allozyme technique is known to detect a low level of polymorphism with regard to other molecular techniques, and underestimate gene variation (Lowe et al. 2004). Genetic diversity values obtained in our study are congruent with a species having open populations and ample distribution with high gene flow probabilities (Roux et al. 2007; Bouzid et al. 2008).

Consequently, the choice of appropriate molecular markers is very important to study genetic variation at the intra-specific level. In the present study, all mainland populations presented high and similar levels of polymorphism and gene diversity coefficients, whereas the island population of Cozumel presented the lowest values. Generally founded from a small number of individuals (founder effect), island populations usually present less genetic diversity than mainland populations and are often inbred (limited gene flow) (Frankham et al. 2005). However, the values of the Cozumel population did not indicate a threatened population and suggest recent colonization of the island, or an ancient colonization with the occasional introduction, most likely by

man, of new individuals from the mainland that can decrease the genetic drift effect.

This study clearly showed the potential of ISSR markers to evaluate genetic diversity in tarantula spiders, and proved an attractive alternative to other molecular markers.

ACKNOWLEDGMENTS

We are grateful to the people of the villages Ley de Fomento, Conhuas, Zoh-Laguna, Luis Echeverría, and Raudales for granting us access to their land and for their hospitality during our stay. We thank Héctor González Cortés of the “Fundación de Parques y Museos de Cozumel” for providing logistic support during our visit to Cozumel. We are grateful to Janneth Adriana Padilla Saldivar of El Colegio de la Frontera Sur (ECOSUR) for producing Figure 1. An earlier version of this manuscript benefited from the insights and comments of Sophie Calmé (ECOSUR) and Peter Winterton.

LITERATURE CITED

- Abbot, P. 2001. Individual and population variation in invertebrates revealed by Inter Simple Sequence Repeats (ISSRs). *Journal of Insect Science* 1:8.
- Arnedo, M.A. & M.A. Ferrández. 2007. Mitochondrial markers reveal deep population subdivision in the European protected spider *Macrothele calpeiana* (Walckenaer, 1805) (Araneae, Hexathelidae). *Conservation Genetics* 8:1147–1162.
- Behura, S.K. 2006. Molecular marker systems in insects: current trends and future avenues. *Molecular Ecology* 15:3087–3113.
- Bond, J.E., D.A. Beamer, T. Lamb & M. Hedin. 2006. Combining genetics and geospatial analyses to infer population extinction in mygalomorph spiders endemic to the Los Angeles region. *Animal Conservation* 9:145–157.
- Bond, J.E., M.C. Hedin, M.G. Ramirez & B.D. Opell. 2001. Deep molecular divergence in the absence of morphological and ecological change in the Californian coastal dune endemic trapdoor spider *Aptostichus simus*. *Molecular Ecology* 10:899–910.
- Bouzid, W., S. Lek, M. Macé, O. Ben Hassine, R. Etienne, L. Legal & G. Loot. 2008. Genetic diversity of *Ligula intestinalis* (Cestoda: Diphyllobothriidea) based on analysis of inter-simple sequence repeat markers. *Journal of Zoological Systematics and Evolutionary Research* 46:289–291.
- Carter, N. 1997. Who's on CITES and why? *Forum of the American Tarantula Society* 6:172–173.
- Casu, M., F. Maltagliati, P. Cossu, T. Lai, M.C. Galletti, A. Castelli & J.A. Commiato. 2005. Fine-grained spatial genetic structure in the bivalve *Gemma gemma* from Maine and Virginia (USA), as revealed by Inter-Simple Sequence Repeat markers. *Journal of Experimental Marine Biology and Ecology* 325:46–54.
- Chatterjee, S.N. & T.P. Mohandas. 2003. Identification of ISSR markers associated with productivity traits in silkworm, *Bombyx mori* L. *Genome* 46:438–447.
- Edwards, G.B. & K.L. Hibbard. 1999. The Mexican Redrump, *Brachypelma vagans* (Araneae: Theraphosidae), an exotic tarantula established in Florida. *Entomology Circular Number 394*, Florida Department of Agriculture & Consumer Services, Division of Plant Industry, Gainesville, Florida. 2 pp.
- Excoffier, L., P.E. Smouse & J.M. Quattro. 1992. Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction sites. *Genetics* 131:479–491.
- Frankham, R., J.D. Ballou & D.A. Briscoe. 2005. *Introduction to Conservation Genetics*. Cambridge University Press, Cambridge, UK. 617 pp.
- Guicking, D., R.A. Griffiths, R.D. Moore, U. Joger & M. Wink. 2006. Introduced alien or persecuted native? Resolving the origin

- of the viperine snake (*Natrix maura*) on Mallorca. *Biodiversity and Conservation* 15:3045–3054.
- Hettle, S.J.H., W. Hall, S.L. Dyas, D.J. Beaumont, D.M. Caplan & D.J. Curtis. 1997. Assessment of the Random Amplified Polymorphic DNA (RAPD) technique for the analysis of spider populations. *Journal of Arachnology* 25:222–227.
- Hoffman, E.A., F.W. Schueler, A.G. Jones & M.S. Blouin. 2006. An analysis of selection on a colour polymorphism in the northern leopard frog. *Molecular Ecology* 15:2627–2641.
- Hundsdoerfer, A.K. & M. Wink. 2006. Incongruence of morphology and genetic markers in *Hyles tithymali* (Lepidoptera: Sphingidae) from the Canary Islands. *Journal of Zoological Systematics and Evolutionary Research* 44:316–322.
- Joger, U., U. Fritz, D. Guicking, S. Kalyabina-Hauf, Z.T. Nagy & M. Wink. 2007. Phylogeography of western Palaearctic reptiles – Spatial and temporal speciation patterns. *Zoologischer Anzeiger* 246:293–313.
- Kothera, L., C.M. Richards & S.E. Carney. 2007. Genetic diversity and structure in the rare Colorado endemic plant *Pisaria bellii* (Brassicaceae). *Conservation Genetics* 8:1043–1050.
- Lewontin, R.C. 1972. The apportionment of human diversity. *Evolutionary Biology* 6:381–398.
- Locht, A., M. Yáñez & I. Vázquez. 1999. Distribution and natural history of Mexican species of *Brachypelma* and *Brachypelmides* (Theraphosidae, Theraphosinae) with morphological evidence for their synonymy. *Journal of Arachnology* 27:196–200.
- Longhorn, S.J. 2002. Non-lethal DNA sampling from CITES II protected ‘tarantula’ spiders of Belize. *Las Cuevas Newsletter* 9:8–9.
- Longhorn, S.J., M. Nicholas, J. Chuter & A.P. Volger. 2007. The utility of molecular markers from non-lethal DNA samples of the CITES II protected “tarantula” *Brachypelma vagans* (Araneae, Theraphosidae). *Journal of Arachnology* 35:278–292.
- Lowe, A., S. Harris & P. Ashton. 2004. *Ecological Genetics: Design, Analysis, and Application*. Blackwell Publishing Company, Oxford, UK. 326 pp.
- Luque, C., L. Legal, H. Staudter, C. Gers & M. Wink. 2002. ISSR (inter simple sequence repeats) as genetic markers in noctuids (Lepidoptera). *Hereditas* 136:251–253.
- Lynch, M. & B.G. Milligan. 1994. Analysis of population genetic structure with RAPD markers. *Molecular Ecology* 3:91–99.
- Machkour-M’Rabet, S., Y. Hénaut, R. Rojo & S. Calmé. 2005. A not so natural history of the tarantula *Brachypelma vagans*: interaction of the human activity. *Journal of Natural History* 39:2515–2523.
- Machkour-M’Rabet, S., Y. Hénaut, A. Sepúlveda, R. Rojo, S. Calmé & V. Geissen. 2007. Soil preference and burrow structure of an endangered tarantula, *Brachypelma vagans* (Mygalomorphae: Theraphosidae). *Journal of Natural History* 41:1025–1033.
- Nagaraju, J., K. Reddy, G. Nagaraja & B. Sethuraman. 2001. Comparison of multilocus RFLPs and PCR-based marker systems for genetic analysis of the silk-worm, *Bombix mori*. *Heredity* 86:588–597.
- Peakall, R. & P.E. Smouse. 2006. GenAlEx 6: Genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* 6:288–295.
- Pedersen, A.A. & V. Loeschcke. 2001. Conservation genetics of peripheral populations of mygalomorph spider *Atypus affinis* (Atypidae) in northern Europe. *Molecular Ecology* 10:1133–1142.
- Ramirez, M.G. & B. Chi. 2004. Cryptic speciation, genetic diversity and gene flow in the California turret spider *Atypoides riversi* (Araneae: Antrodiaetidae). *Biological Journal of the Linnean Society* 82:27–37.
- Ramirez, M.G. & J.L. Froehlig. 1997. Minimal genetic variation in a coastal dune arthropod: the trapdoor spider *Aptostichus simus* (Cyrtachenidae). *Conservation Biology* 11:256–259.
- Reichling, S.B. 2000. Group dispersal in juvenile *Brachypelma vagans* (Araneae, Theraphosidae). *Journal of Arachnology* 28:248–250.
- Roux, O., M. Gevrey, L. Arvanitakis, C. Gers, D. Bordat & L. Legal. 2007. ISSR-PCR: Tool for discrimination and genetic structure analysis of *Plutella xylostella* populations native to different geographical areas. *Molecular Phylogenetics and Evolution* 43:240–250.
- Smith, A. 1994. *Tarantula Spiders: Tarantulas of the USA and Mexico*. Fitzgerald Publishing, London. 200 pp.
- Wink, M. 2006. Use of DNA markers to study bird migration. *Journal of Ornithology* 147:234–244.
- Wink, M., H. Sauer-Gurth & E. Gwinner. 2002. A molecular phylogeny of stonechats and related turdids inferred from mitochondrial DNA sequences and genomic fingerprinting by ISSR-PCR. *British Birds* 95:349–355.
- Wolfe, A.D. & A. Liston. 1998. Contributions of PCR-based methods to plant systematics and evolutionary biology. Pp. 43–86. *In: Plant Molecular Systematics II* (D.E. Soltis, P.S. Soltis & J.J. Doyle, eds.). Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Yáñez, M. & G. Floater. 2000. Spatial distribution and habitat preference of the endangered tarantula, *Brachypelma klaasi* (Araneae: Theraphosidae) in Mexico. *Biodiversity and Conservation* 9:795–810.
- Yáñez, M., A. Locht & R. Macias-Ordóñez. 1999. Courtship and mating behavior of *Brachypelma klaasi* (Araneae, Theraphosidae). *Journal of Arachnology* 27:165–170.
- Yeh, F.C., R.C. Yang, T.B.J. Boyle, Z.H. Ye & J.X. Mao. 1997. POPGENE, the User-Friendly Shareware for Population Genetic Analysis. Version 1.21. Molecular Biology and Biotechnology Center, University of Alberta, Canada. Online at <http://www.ualberta.ca/~fyeh>
- Zietkiewicz, E., A. Rafalski & D. Labuda. 1994. Genome fingerprinting by simple sequence repeat (SSR)-anchored polymerase chain reaction amplification. *Genomics* 20:176–183.

Manuscript received 18 March 2008, revised 10 October 2008.