

## Impact of elevated CO<sub>2</sub> on growth, development, and reproduction of the wolf spider, *Pardosa astrigera* (Araneae: Lycosidae)

Ling Zuo<sup>1</sup>, Cheng Chen<sup>1</sup>, Ling Qi<sup>1</sup>, Feng-Xiang Liu<sup>1</sup>, Yue-li Yun<sup>1</sup> and Yu Peng<sup>1,2</sup>: <sup>1</sup>Centre for Behavioral Ecology & Evolution, College of Life Sciences, Hubei University, Wuhan 430062, P.R. China; <sup>2</sup>Hubei Collaborative Innovation Center for Green Transformation of Bio-Resources, College of Life Sciences, Hubei University, Wuhan 430062, P.R. China. E-mail: pengyu@hubu.edu.cn

**Abstract.** The effects of elevated CO<sub>2</sub> concentration on spiders were studied using third-instar spiderlings of the wolf spider *Pardosa astrigera* L. Koch 1878 in CO<sub>2</sub> climate chambers with two different concentrations of CO<sub>2</sub> (low, 370 ppm and high, 750 ppm). The food intake and total developmental period of spiderlings reared at high CO<sub>2</sub> concentration increased significantly, and the body length and weight of adult spiders decreased compared to those in the low-CO<sub>2</sub> group. The oviposition rate of female *P. astrigera* and the hatching rate of eggs did not differ between the high- and low-CO<sub>2</sub> groups, but the number of egg sacs and the total number of eggs produced by females from the high-CO<sub>2</sub> group decreased. These results suggest that elevated CO<sub>2</sub> concentrations are harmful to the growth, development, and reproduction of *P. astrigera*.

**Keywords:** Body length, body weight, oviposition rate

Elevated atmospheric carbon dioxide (CO<sub>2</sub>) concentration is thought to be the main reason for global climate change and the greenhouse effect. Climatic records show that the atmospheric CO<sub>2</sub> concentration has increased at an annual rate of 1.9 ppm since 1995. It is estimated that the atmospheric CO<sub>2</sub> concentration will rise to 540–970 ppm by 2100 (IPCC 2013). Because of the increasing atmospheric CO<sub>2</sub> concentration, studying its effects on organisms has been a very popular subject in ecological research (Cannon 1998; Stiling & Cornelissen 2007; Ge et al. 2010; Cornelissen 2011; Foss et al. 2013). Studies to date have mostly investigated the influence of elevated CO<sub>2</sub> on plants, phytophagous insects, and natural enemy insects (predators and parasitoids) (Lincoln et al. 1984; Bezemer & Jones 1998; Wu et al. 2006; Chen et al. 2007; Ge et al. 2010). However, it remains unclear how an elevated CO<sub>2</sub> concentration impacts spiders, a key predator of insects.

Elevated atmospheric CO<sub>2</sub> concentrations have direct effects on the growth, chemistry, and physiology of plants (Lindroth et al. 1993; Roth & Lindroth 1995; Lin & Wang 2002; Veteli et al. 2002; Chen et al. 2005a). For phytophagous and natural enemy insects, elevated CO<sub>2</sub> concentration indirectly affect their growth, development, and reproduction (Brooks & Whittaker 1998; Deng et al. 2002; Chen et al. 2005b, 2007; Wu et al. 2006, 2007; Cornelissen 2011; Foss et al. 2013).

Spiders play important roles in controlling pests (Marc et al. 1999; Barrion et al. 2012) and monitoring environmental conditions (Babczynska et al. 2011; Chen et al. 2011). Understanding the effects of elevated CO<sub>2</sub> concentration on spiders will contribute to better protection and utilization of them. The objective of this study was to determine the effects of elevated CO<sub>2</sub> concentration on spiders. We investigated the food intake, molting, clutch size and egg hatch of the wolf spider *Pardosa astrigera* L. Koch 1878 under different CO<sub>2</sub> concentrations. Based on previous studies about predatory insects (Chen et al. 2007; Ge et al. 2010), we hypothesized that an elevated CO<sub>2</sub> concentration could have either harmful or beneficial effects on the spiders.

## METHODS

**Spider collection and rearing.**—Subadult individuals of *P. astrigera* were collected from farm fields in Ma'anshan Forest Park, Wuhan (30° 52'N, 114° 31'E), Hubei Province, China, in April and May 2012. Voucher specimens were deposited in the Centre for Behavioral Ecology & Evolution, College of Life Sciences, Hubei University, China.

Spiders were kept individually in cylindrical glass tubes (diameter 2 cm, height 12 cm) with a layer of sponge (1.5 cm thick) moistened with water on the bottom. The tubes were plugged with cotton. The spiders were kept in a chamber at 24°C and relative humidity of 60–80% under a light: dark cycle of 14:10 h (lights turned on at 08:00). Every two days, we fed the spiders with adults of *Drosophila melanogaster* cultured under normal CO<sub>2</sub> concentration. Two days post-maturation, females and males were placed together for mating. The male was removed after the female deposited the first egg sac. Following the emergence of second-instar spiderlings and dispersal from the female's abdomen, siblings from different egg sacs were separated and reared individually, with *D. melanogaster* provided as food. Once the second-instar spiderlings molted, they were used for the following experiments.

**Spiderling rearing in different CO<sub>2</sub> concentrations.**—A total of 300 third-instar spiderlings were randomly selected from different egg sacs and reared individually in cylindrical glass tubes (diameter 2 cm, height 12 cm) and then divided into two groups (150 in each group). The spiderlings were reared under low CO<sub>2</sub> (370 ppm) and high CO<sub>2</sub> (750 ppm) concentrations in CO<sub>2</sub> artificial climate chambers (CC350TLHC type, Changzhou Okefenokee Instrument Co., LTD). The concentrations of CO<sub>2</sub> were set according to Chen et al. (2005b) and Ge et al. (2010); the current concentration of atmospheric CO<sub>2</sub> is 370 ppm, and 750 ppm is approximately double the current level. Each CO<sub>2</sub> concentration was repeated three times by using three chambers, with 50 spiderlings in each chamber. The chambers were set at 25°C and relative humidity of 40%–60% with a light: dark cycle of 14:10 h (lights turned on at




Figure 1.—Effect of different CO<sub>2</sub> concentrations on food intake of *Pardosa astrigera* fed on *Drosophila melanogaster* ( $n = 10$ ). Different letters above the columns indicate significant differences in food intake of spiders maintained under different CO<sub>2</sub> concentrations (t-test,  $df = 9$ ,  $t = 3.39$ ,  $P = 0.008$ ).

08:00). The spiderlings were fed adults of *D. melanogaster* and the sponges were replaced every two days.

**Measurement of food intake of spiderlings.**—The food intake of third-instar spiderlings ( $n = 10$ ) under different CO<sub>2</sub> concentrations was investigated. Fifteen fruit flies (*D. melanogaster*) were provided at 17:00 and the number of living flies was recorded the next day at 09:00. The difference between the initial and final number of flies was taken to be the food intake of the spiderlings.

**Assessment of developmental duration, body length, weight, and fecundity.**—Molts were recorded when exuviae were observed in the tube, and the time between molts was used as a measure of developmental duration. Two days post-maturation, the females of *P. astrigera* ( $n = 15$ ) were randomly paired with a mature male and left together for 48 h to mate. Oviposition rate, the number of egg sacs laid by the females ( $n = 10$ ), the total number of eggs ( $n = 15$ ), and hatching rate of the eggs ( $n = 4$ ) were observed and recorded for each treatment (Chen et al. 2011). Ten female and 10 male spiders from different treatments were randomly selected and weighed to the nearest 0.01 mg by using electronic balance (FA1004N type, HANGPING). Then, the above 20 spider individuals were killed in 75% alcohol to measure the body

Table 1.—Developmental durations (days) of spiderlings of *Pardosa astrigera* under different CO<sub>2</sub> concentrations. Data were expressed as Mean ± standard deviation. The differences between different CO<sub>2</sub> concentrations were compared by Student’s t-test. The same notation is used in Tables 2– 4.

Items	CO <sub>2</sub> concentration (ppm)		<i>t</i>	<i>P</i>
	370	750		
3rd instar	15.4 ± 0.6	20.6 ± 0.9	2.31	0.025
4th instar	13.2 ± 0.5	19.0 ± 0.7	2.29	0.031
5th instar	16.3 ± 0.9	25.1 ± 0.4	2.97	0.008
6th instar	17.1 ± 0.9	22.4 ± 0.4	2.58	0.022
Total	61.8 ± 2.4	87.1 ± 1.5	2.96	0.0037

Table 2.—Body lengths (mm) of adult *Pardosa astrigera* under different CO<sub>2</sub> concentrations ( $n = 10$ ).

Sex of spider	CO <sub>2</sub> concentration (ppm)		<i>t</i>	<i>P</i>
	370	750		
Female	9.70 ± 0.84	7.10 ± 0.42	2.66	0.026
Male	8.30 ± 0.45	5.40 ± 0.42	3.14	0.012

lengths (from front edge of carapace to the end of abdomen) by using an ocular micrometer under a microscope (DFC495 type, LEICA).

**Statistical analysis.**—Data were expressed as mean ± standard deviation. The differences between treatments were compared by Student’s t-test.

RESULTS

The consumption of *D. melanogaster* by the third-instar spiderlings of *P. astrigera* reared at the high CO<sub>2</sub> concentration was significantly higher than at the low CO<sub>2</sub> concentration (t-test,  $df = 9$ ,  $t = 3.39$ ,  $P = 0.008$ ) (Figure 1).

The duration of development (days) for each instar (t-test,  $P < 0.05$ ) and the total duration (t-test,  $df1 = 65$ ,  $df2 = 53$ ,  $t = 2.96$ ,  $P = 0.0037$ ) for the spiderlings were significantly longer for the spiderlings kept under high CO<sub>2</sub> concentration than under low CO<sub>2</sub> concentration (Table 1). This suggested that elevated CO<sub>2</sub> caused the spiderlings to mature later, prolonging the total development time.

The body length of female *P. astrigera* was longer than in males kept under the same CO<sub>2</sub> concentration (Table 2). Body lengths of both adult females (t-test,  $df = 9$ ,  $t = 2.66$ ,  $P = 0.026$ ) and males (t-test,  $df = 9$ ,  $t = 3.14$ ,  $P = 0.012$ ) of the high-CO<sub>2</sub> group were shorter than those of the low-CO<sub>2</sub> group. This showed that the body lengths of mature spiders were decreased under the high CO<sub>2</sub> concentration.

The body weight of females of *P. astrigera* was also higher than that of males under the same CO<sub>2</sub> concentration. The body weights of both female (t-test,  $df = 9$ ,  $t = 4.78$ ,  $P = 0.001$ ) and male spiders (t-test,  $df = 9$ ,  $t = 2.92$ ,  $P = 0.017$ ) reared at the low CO<sub>2</sub> concentration were heavier than those reared at high CO<sub>2</sub> (Table 3).

There were no differences in the oviposition rate of female *P. astrigera* (t-test,  $df = 14$ ,  $t = 0$ ,  $P = 1.0$ ) or the hatching rate of eggs (t-test,  $df = 3$ ,  $t = 1.01$ ,  $P = 0.37$ ) between the high- and low-CO<sub>2</sub> groups (Table 4). However, the number of egg sacs produced by the females from the high-CO<sub>2</sub> group was lower than the number produced by the low-CO<sub>2</sub> group (t-test,  $df = 9$ ,  $t = 2.69$ ,  $P = 0.025$ ), and the total number of eggs of the high-CO<sub>2</sub> group was also lower than in the low-CO<sub>2</sub> group (t-test,  $df = 14$ ,  $t = 3.49$ ,  $P = 0.0036$ ).

Table 3.—Body masses (mg) of adult *Pardosa astrigera* under different CO<sub>2</sub> concentrations ( $n = 10$ ).

Sex of spider	CO <sub>2</sub> concentration (ppm)		<i>t</i>	<i>P</i>
	370	750		
Female	41.3 ± 8.6	33.5 ± 2.7	4.78	0.001
Male	31.9 ± 0.6	22.4 ± 0.5	2.92	0.017

Table 4.—Fecundity of female *Pardosa astrigera* under different CO<sub>2</sub> concentrations

Items	CO <sub>2</sub> concentration (ppm)		<i>t</i>	<i>P</i>
	370	750		
Oviposition rate (%)	100 ± 0	100 ± 0	0	1.0
Number of egg sacs	5.6 ± 0.9	2.8 ± 0.8	2.69	0.025
Total number of eggs	229.6 ± 39.8	125.3 ± 34.9	3.49	0.0036
Hatching rate (%)	100 ± 0	83.7 ± 15.6	1.01	0.37

## DISCUSSION

Under a high concentration of atmospheric CO<sub>2</sub>, the nitrogen content of plants decreases, and defoliating insects will increase their feeding rate in order to compensate for the nutritional loss caused by the lower nitrogen content of the host plant (Lincoln et al. 1984). Chen et al. (2007) found that the larvae of lady beetles (*Harmonia axyridis*) consumed more cotton aphids *Aphis gossypii* fed on cotton plants grown in elevated CO<sub>2</sub> in order to compensate for the reduced soluble protein in *A. gossypii* owing to the decrease in foliar N and the increase in the C:N ratio in the cotton plants. Our results showed that the third-instar spiderlings of *P. astrigera* also consumed more fruit flies at the high than at the low CO<sub>2</sub> concentration. This phenomenon may have been caused by the larger amounts of energy required by the spiders and higher respiration rates (Foss et al. 2013) to compensate for their increased activity in the high-CO<sub>2</sub> environment.

High atmospheric CO<sub>2</sub> concentration could affect enzyme activity in the cotton bollworm *Helicoverpa armigera*, and the available nutrients would decline significantly, thereby impairing the growth and development of *H. armigera* (Chen et al. 2005b). The body length and weight similarly decreased in *H. armigera* (Chen et al. 2005b). When second-stadium gypsy moth (*Lymantria dispar*) larvae were grown under high CO<sub>2</sub> concentration, their body weight was lower than in the normal CO<sub>2</sub> concentration (Wang et al. 2006). Our former results showed that when *P. astrigera* was subjected to the stress of Pb or Zn, its developmental duration lengthened (Chen et al. 2011). Similar results were found in our present study. The development time of *P. astrigera* lengthened and the body length and weight of mature spiders decreased. This suggests that these different environmental factors can cause developmental delay in this spider species.

Our results indicated that the fecundity of *P. astrigera* was significantly decreased at the high CO<sub>2</sub> concentration. Brooks & Whittaker (1998) found that the number of eggs produced by females of the green dock beetle *Gastrophysa viridula* that were continuously cultured for three generations, decreased under a high atmospheric CO<sub>2</sub> concentration. At the high atmospheric CO<sub>2</sub> concentration, the number of eggs laid by females of *H. armigera* was markedly reduced (Chen et al. 2005b). However, some studies have had different outcomes. Bezemer & Jones (1998) showed that the number of eggs laid by females of the winter moth *Operophtera brumata* could increase, and the number of eggs laid by *G. viridula* did not differ significantly between high and normal CO<sub>2</sub> concentrations (Brooks & Whittaker 1998). In the present study, female spiders laid fewer eggs under the stress of high CO<sub>2</sub> concentration. Taken together, these studies show that the

impact of elevated CO<sub>2</sub> on the reproduction of arthropods varies among species.

In summary, the high CO<sub>2</sub> concentration proved to be harmful to the growth, development, and reproduction of *P. astrigera*. The results of this experiment increase understanding of the responses of spiders reared in elevated atmospheric CO<sub>2</sub> concentration.

## ACKNOWLEDGMENTS

This study was supported by the National Natural Science Fund of China (31071895), and the Key Scientific and Technological Projects of Hubei (2013BEC035) and Wuhan (2014020101010074). We are grateful to two anonymous reviewers for suggestions and improvements to this manuscript. We also thank Dr. Janet W. Reid (JWR Associates, New York, USA) for editing the English text.

## LITERATURE CITED

- Babczyńska, A., G. Wilczek, E. Szulińska & I. Franiel. 2011. Quantitative immunodetection of metallothioneins in relation to metals concentration in spiders from variously polluted areas. *Ecotoxicology and Environmental Safety* 74:1498–1503.
- Barrion, A.T., S.S. Villareal, J.L.A. Catindig, D. Cai, Q.H. Yuan & K.L. Heong. 2012. The spider fauna in the rice agricultural landscape of Hainan Island, China: composition, abundance and feeding structure. *Asia Life Sciences* 21:625–651.
- Bezemer, T.M. & T.H. Jones. 1998. Plant-insect herbivore interactions in elevated atmospheric CO<sub>2</sub>: quantitative analysis and guild effects. *Oikos* 82:212–222.
- Brooks, G.L. & J.B. Whittaker. 1998. Responses of multiple generations of *Gastrophysa viridula*, feeding on *Rumex obtusifolius*, to elevated CO<sub>2</sub>. *Global Change Biology* 4:63–75.
- Cannon, R.J.C. 1998. The implications of predicted climate change for insect pests in the UK, with emphasis on non-indigenous species. *Global Change Biology* 4:785–796.
- Chen, F.J., G. Wu, F. Ge, M.N. Parajulee & R.B. Shrestha. 2005a. Effects of elevated CO<sub>2</sub> and transgenic Bt cotton on plant chemistry, performance, and feeding of an insect herbivore, the cotton bollworm. *Entomologia Experimentalis et Applicata* 115:341–350.
- Chen, F.J., G. Wu, J. Lu & F. Ge. 2005b. Effects of elevated CO<sub>2</sub> on the foraging behavior of cotton bollworm, *Helicoverpa armigera*. *Insect Science* 12:359–365.
- Chen, F.J., G. Wu, M.N. Parajulee & F. Ge. 2007. Impact of elevated CO<sub>2</sub> on the third trophic level: a predator *Harmonia axyridis* and a parasitoid *Aphidius picipes*. *Biocontrol Science and Technology* 17:313–324.
- Chen, X.Q., Z.T. Zhang, R. Liu, X.L. Zhang, J. Chen & Y. Peng. 2011. Effects of the metals lead and zinc on the growth, development, and reproduction of *Pardosa astrigera* (Araneae: Lycosidae). *Bulletin of Environmental Contamination and Toxicology* 86:203–207.
- Cornelissen, T. 2011. Climate change and its effects on terrestrial insects and herbivory patterns. *Neotropical Entomology* 40:155–63.
- Deng, Y.X., Z.M. Zhao & L.S. Li. 2002. Control of high CO<sub>2</sub> concentrations on *Rhizopertha dominica* and *Tribolium confusum*. *Grain Storage* 31:43–56.
- Foss, A.R., W.J. Mattson & T.M. Trier. 2013. Effects of elevated CO<sub>2</sub> leaf diets on gypsy moth (Lepidoptera: Lymantriidae) respiration rates. *Environmental Entomology* 2:503–514.
- Ge, F., F.J. Chen, G. Wu & Y.C. Sun. 2010. Research advance on the response of insects to elevated CO<sub>2</sub> in China. *Chinese Bulletin of Entomology* 47:229–235.

- IPCC. 2013. Summary for policymakers. Pp. 3–29. *In* Climate Change 2013: The Physical Science Basis Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. (T.F. Stocker, D. Qin, G.K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex & P.M. Midgley, eds.). Cambridge University Press, Cambridge.
- Lin, J.S. & G.X. Wang. 2002. Double CO<sub>2</sub> could improve the drought tolerance better in sensitive cultivars than in tolerant cultivars in spring wheat. *Plant Science* 163:627–637.
- Lincoln, D.E., N. Sionit & B.R. Strain. 1984. Growth and feeding response of *Pseudoplusia includens* (Lepidoptera: Noctuidae) to host plant growth in controlled carbon dioxide atmosphere. *Environmental Entomology* 13:1527–1530.
- Lindroth, R.L., K.K. Kinney & C.L. Platz. 1993. Responses of deciduous trees to elevated atmospheric CO<sub>2</sub>: productivity, phytochemistry, and insect performance. *Ecology* 74:763–777.
- Marc, P., A. Canard & F. Ysnel. 1999. Spiders (Araneae) useful for pest limitation and bioindication. *Agriculture, Ecosystems & Environment* 74:229–273.
- Roth, S.K. & R.L. Lindroth. 1995. Elevated atmospheric CO<sub>2</sub>: effects on phytochemistry, insect performance and insect-parasitoid interactions. *Global Change Biology* 1:173–182.
- Stiling, P. & T. Cornelissen. 2007. How does elevated carbon dioxide (CO<sub>2</sub>) affect plant–herbivore interactions? A field experiment and meta-analysis of CO<sub>2</sub>-mediated changes on plant chemistry and herbivore performance. *Global Change Biology* 13:1–20.
- Veteli, T.O., K. Kuokkanen, R. Julkunen-Tiitto, H. Roininen & J. Tahvanainen. 2002. Effects of elevated CO<sub>2</sub> and temperature on plant growth and herbivore defensive chemistry. *Global Change Biology* 8:1240–1252.
- Wang, X.W., L.Z. Ji & Y. Liu. 2006. Effects of increased atmospheric CO<sub>2</sub> on nutritional contents in poplar tissues and larval growth of gypsy moth. *Acta Ecologica Sinica* 10:3166–3174.
- Wu, G., F.J. Chen & F. Ge. 2006. Response of multiple generations of cotton bollworm *Helicoverpa armigera* Hubner, feeding on spring wheat, to elevated CO<sub>2</sub>. *Journal of Applied Entomology* 130:2–9.
- Wu, G., F.J. Chen & F. Ge. 2007. Transgenic *Bacillus thuringiensis* (Bt) cotton (*Gossypium hirsutum*) allomone response to cotton aphid, *Aphis gossypii* in a closed-dynamics CO<sub>2</sub> chamber (CDCC). *Journal of Plant Research* 120:679–685.

*Manuscript received 3 May 2014, revised 1 December 2014.*