

THE FIRST DNA PHYLOGENY OF FOUR SPECIES OF *MESOBUTHUS* (SCORPIONES, BUTHIDAE) FROM EURASIA

Benjamin Gantenbein: Institute of Cell, Animal and Population Biology, University of Edinburgh, West Mains Road, Edinburgh EH9 3JT, United Kingdom

Victor Fet: Department of Biological Sciences, Marshall University, Huntington, West Virginia 25755-2510 USA

Alexander V. Gromov: Institute of Zoology, Akademgorodok, Almaty, 480060, Republic of Kazakhstan

ABSTRACT. The first molecular phylogeny is presented for four species of the scorpion genus *Mesobuthus*, based on DNA sequences of three gene fragments (two mitochondrial and one nuclear protein coding gene, ~1 kb). The inferred phylogeny based on a pooled maximum likelihood analysis revealed a clear deep splitting between the “western clade” consisting of *M. gibbosus* and *M. cyprius* (Greece/Anatolia, Cyprus) and the “eastern clade” consisting of *M. eupeus* and *M. caucasicus* (Anatolia/Central Asia). The species *M. caucasicus* (recently placed in the genus *Olivierus* Farzanpay 1987) groups monophyletically within *Mesobuthus*; thus, the genus *Olivierus* is synonymized here with *Mesobuthus*. Sequences of *M. eupeus* and *M. caucasicus* sampled mainly from Kazakhstan and Uzbekistan are highly structured, indicating the possible existence of multiple species.

Keywords: Scorpions, Buthidae, *Mesobuthus*, phylogeny, DNA, 16S, *coxI*, protein kinase, biogeography

The genus *Mesobuthus* Vachon 1950 (Scorpiones, Buthidae) currently includes 12 species (Fet & Lowe 2000; Gantenbein et al. 2000b); its type species is *Mesobuthus eupeus* (C.L. Koch 1839). Except for *M. gibbosus* (Brullé 1832), which is found in the Balkans and Turkey, the diversity of this genus is concentrated in Asia. Numerous species and subspecies are distributed from Turkey to Korea, with the centers of diversity in Central Asia and Iran. *Mesobuthus* species are the most common and abundant scorpions in a variety of arid habitats, from sand deserts to high mountains over 3000 m (Fet 1989, 1994). They are found up to 50°N in Kazakhstan, the northern limit of the natural range of the Old World scorpions (Gromov 2001).

Although Buthidae are the most diverse and medically important family of scorpions (Fet & Lowe 2000), there has been no attempt so far to produce a phylogenetic analysis of this family. Especially powerful are phylogenies based on DNA sequence data in combination with morphology (Gantenbein et al. 1999a, 2000a; Fet et al. 2001). Our first applications of this technique in Buthidae refer to the gen-

era *Buthus* Leach 1815 (Gantenbein et al. 1999b; Gantenbein & Largiadèr 2003) and *Centruroides* Marx 1890 (Gantenbein et al. 2001; Towler et al. 2001); we also published a pilot phylogeny of 17 buthid genera (Fet et al. 2003). Molecular markers helped to define island species, where neither the biological species concept nor any other species concept can be applied (Gantenbein et al. 2000b, 2001).

METHODS

The currently studied available material belonged to four species: *Mesobuthus gibbosus* (Greece, Turkey), *M. cyprius* Gantenbein & Kropf 2000 (Cyprus), *M. eupeus* (C.L. Koch 1839) (Turkey, Kazakhstan, Turkmenistan, Uzbekistan, China), and *M. caucasicus* (Nordmann 1840) (Kazakhstan, Turkmenistan, Uzbekistan) (see Table 1 for locality information). For DNA analyses, the total DNA was extracted from fresh or preserved (94–98% ethanol) muscle tissue using a standard phenol/chloroform and precipitation method (Sambrook et al. 1989). We amplified a ca. 450 base pair (bp) fragment of the 16S rRNA mitochondrial (mt) DNA by polymerase chain reaction (PCR) using the primers and condi-

Table 1.—Sampling sites and country of origin of *Mesobuthus* species used in this study.

| Abbreviation | Country | Locality | Coll. | EMBL accession nos. | | |
|---|--------------|-----------------------------|-----------------------------------|---------------------|-------------|-------------|
| | | | | 16S | <i>coxI</i> | PK |
| <i>M. caucasicus</i> (Nordmann 1840) | | | | | | |
| <i>McaKZa1</i> | Kazakhstan | Kapchagai | Ch. Tarabaev, 1990 | AJ550674 | AJ550692 | AJ550713 |
| <i>McaKZb1</i> | Kazakhstan | Baigakum | V. Fet & A. Gromov, 25-V-2002 | AJ550675 | AJ550693 | AJ550714 |
| <i>McaUZa1</i> | Uzbekistan | Bukhara | V. Fet & A. Gromov, 20-IV-2002 | AJ550676 | AJ550694 | AJ550715 |
| <i>McaUZb1</i> | Uzbekistan | Jarkurgan | V. Fet & A. Gromov, 26-IV-2002 | AJ550677 | AJ550695 | AJ550716 |
| <i>McaUZc1</i> | Uzbekistan | Karakalpak Steppe, Ferghana | V. Fet & A. Gromov, 20-V-2002 | AJ550678 | AJ550696 | AJ550717 |
| <i>McaUZd1</i> | Uzbekistan | Babatag Mountains | V. Fet & A. Gromov, 30-IV-2002 | AJ550679 | AJ550697 | AJ550718 |
| <i>M. cyprius</i> Gantenbein & Kropf 2000 | | | | | | |
| <i>McyCYa1</i> | Cyprus | Tepebasi | A. Scholl, 27-IX-1997 | AJ550680 | AJ550698 | AJ550719 |
| <i>McyCYb1</i> | Cyprus | Kantara | A. Scholl, 20-V-1998 | AJ550681 | AJ550699 | AJ550720 |
| <i>M. eupeus mongolicus</i> (Birula 1911) | | | | | | |
| <i>MeuGobi1</i> | China | Gobi Desert | A. Davidson, 30-VII-1998 | AJ550682 | AJ550700 | AJ550721 |
| <i>M. eupeus eupeus</i> (C.L. Koch 1839) | | | | | | |
| <i>MeuTRa1</i> | Turkey | Gölşehir, Central Anatolia | A. Scholl, 28-V-1998 | AJ550688 | AJ550701 | AJ550722 |
| <i>MeuTRb1-2</i> | Turkey | Cemilköy, Central Anatolia | A. Scholl, 31-V-1998 | AJ550689–90 | AJ550702–03 | AJ550723–24 |
| <i>M. eupeus thersites</i> (C.L. Koch 1839) | | | | | | |
| <i>MeuKZa1</i> | Kazakhstan | Bakanas | A. Gromov, 2-5-VI-2000 | AY228141 | AJ550704 | AJ550725 |
| <i>MeuKZb1</i> | Kazakhstan | Baigakum | V. Fet & A. Gromov, 23-V-2002 | AJ550684 | AJ550705 | AJ550726 |
| <i>MeuKZc2</i> | Kazakhstan | Karatau Mountains | V. Fet & A. Gromov, 27-V-2002 | AJ550685 | AJ550706 | AJ550727 |
| <i>MeuTU1a</i> | Turkmenistan | Repetek, Karakum | V. Fet & A. Gromov, 15-18-IV-2002 | AJ550686 | AJ550707 | AJ550728 |
| <i>MeuUZa1</i> | Uzbekistan | Zarafshan, Kizylkum | A. Gromov, 18-IV-1998 | AJ550687 | AJ550708 | AJ550729 |

Table 1.—Continued.

| Abbreviation | Country | Locality | Coll. | EMBL accession nos. | | |
|--|------------|--------------------------|---------------------------------|---------------------|-------------|----------|
| | | | | 16S | <i>coxI</i> | PK |
| <i>MeuUZb1</i> | Uzbekistan | Babatag Mountains | V. Fet & A. Gromov, 29-IV-2002 | AJ550683 | AJ550709 | AJ550730 |
| <i>Mesobuthus gibbosus</i> (Brullé 1832) | | | | | | |
| <i>MgiGRa1</i> | Greece | Mathia, Peloponnesos | I. & B. Gantenbein, 18-III-1998 | AJ402571 | AJ550710 | AJ550731 |
| <i>MgiGRb1</i> | Greece | Igoumenitsa, Epirus | I. & B. Gantenbein, 28-IV-1998 | AJ550691 | AJ550711 | AJ550732 |
| <i>MgiTRa1</i> | Turkey | Avanos, Central Anatolia | A. Scholl, 28-V-1998 | AJ402587 | AJ550712 | AJ550733 |
| Outgroup: <i>Androctonus australis</i> (Linnaeus 1758) | | | | | | |
| <i>AanTNa1</i> | Tunisia | Nefta | A. Scholl, 27-IV-1999 | AJ506868 | AJ506919 | AJ550734 |

tions of Gantenbein et al. (1999). For the partial amplification of the cytochrome oxidase I (*coxI*) gene we used the primers *LCO* (Folmer et al. 1994) and *Nancy* (Simon et al. 1994) which amplify a ~850 bp fragment (positions ~30–850 of the *Drosophila yakuba* sequence; Flybase: FBgn0013179). We used the following PCR profile: initial denaturation 94 °C for 4 min followed by 40 cycles of 25 s at 94 °C, 20 s at 51 °C, and 90 s at 72 °C. In addition, we designed new buthid-specific PCR primers from a clone (03B09) of an EST library of *Mesobuthus gibbosus* (unpublished data, Gantenbein et al., in preparation) to amplify a ~360 bp fragment of the nuclear protein kinase (PK) gene (Flybase locus CG11221, identified from BLASTX against the *Drosophila* protein database, similarity 43%, Expectation = 5e-14). The primers were *03B09for* 5'-TCT GAT GTA TGG CAG ATG GCA ATG-3' and *03B09rev* 5'-CGA ACT

CAA GAT CCA CTC CTG TAC TCG-3'. We used the same PCR profile as for *coxI*. PCR primers were removed by polyethylene glycol (PEG 8000) precipitation, and templates were directly sequenced on one strand using one of the PCR primers and DYEnamic ET Dye Terminator Kit (Amersham Biosciences). For *coxI* and PK we used the forward PCR-primers for sequencing, and for the 16S we used the same primer as in Gantenbein et al. (1999). Sequencing reactions were ethanol/sodium-acetate precipitated and run on an ABI377XL sequencer (Applied Biosystems, Foster City, CA). All sequences were checked manually for sequencing errors. All sequences were deposited in the EMBL nucleotide sequence database (<http://www.ebi.ac.uk>). As outgroup, we used the Old World buthid *Androctonus australis* (L. 1758) from Nefta, Tunisia (AauTNa1), which is a suitable outgroup for the genus *Mesobuthus* as indicated from

→

Figure 1.—Maximum Likelihood (ML) tree of *Mesobuthus* species from southern Europe, Western and Central Asia inferred from three combined DNA sequence fragments of the mitochondrial 16S, *coxI* and the nuclear PK regions, 1,095 bp (-ln Likelihood was 5505.86). The DNA substitution model was TRN + Γ + I (Tamura & Nei 1993); base frequencies: $\pi_A = 0.29$, $\pi_T = 0.37$, $\pi_C = 0.14$, $R_{matrix} = (A-G = 6.70, A-C = A-T = G-T = 1, C-T = 3.56)$, gamma shape parameter $\alpha = 0.53$, and proportion of invariable sites = 0.37, respectively. The tree was rooted using the outgroup species *Androctonus australis* (AauTNa1). Numbers at nodes refer to bootstrap support given as percentage from 1,000 pseudo replicates by neighbour-joining of ML distances.




Table 2.—Estimates of average rates of synonymous substitutions (K_S) (lower left) and non-synonymous (K_A) (upper right) substitution per site among four species of *Mesobuthus* of two protein-coding gene fragments. Diagonals show average K_S and K_A (in parentheses) among haplotypes within species. Estimates of nucleotide diversity π for each species (Nei & Li 1979) are given in the last column.

| | <i>cyprius</i> | <i>caucasicus</i> | <i>eupeus</i> | <i>gibbosus</i> | diversity π |
|-------------------|----------------|-------------------|---------------|-----------------|-----------------|
| coxI locus | | | | | |
| <i>cyprius</i> | 0.11 (0.009) | 0.297 | 0.024 | 0.006 | 0.037 |
| <i>caucasicus</i> | 0.844 | 0.482 (0.030) | 0.034 | 0.023 | 0.098 |
| <i>eupeus</i> | 0.887 | 0.753 | 0.377 (0.026) | 0.023 | 0.089 |
| <i>gibbosus</i> | 0.538 | 0.892 | 0.692 | 0.223 (0.004) | 0.048 |
| PK locus | | | | | |
| <i>cyprius</i> | 0.038 (0.018) | 0.106 | 0.013 | 0.013 | 0.032 |
| <i>caucasicus</i> | 0.093 | 0.022 (0.000) | 0.004 | 0.004 | 0.008 |
| <i>eupeus</i> | 0.073 | 0.032 | 0.008 (0.000) | 0.009 | 0.002 |
| <i>gibbosus</i> | 0.04 | 0.082 | 0.064 | 0.037 (0.009) | 0.017 |

morphology (Vachon 1952). Voucher specimens will be deposited in the Natural History Museum, Bern, Switzerland (NHMBE).

The resulting 22 16S DNA sequences were aligned using ClustalX (Thompson et al. 1997) and by eye. We applied Maximum Likelihood (ML) to the DNA sequence data. The alignment was 344 bp long. The alignment was deposited in the EMBL nucleotide database (ALIGN_000522). Ambiguities and gaps were stripped out (Swofford et al. 1996), leaving 302 bp. The alignment of the *coxI* and the PK fragments was unambiguous because of the open reading frames (ORF). In order to choose the most appropriate DNA model of nucleotide substitution, we calculated hierarchic likelihood ratio test statistics using MODELTEST 3.06 (Posada & Crandall 1998; Huelsenbeck & Rannala 1997) which is implemented in PAUP* 4.0b10 (Swofford 1998) and calculates the hierarchical likelihood ratio statistics (LRT) of 56 different substitution models based on a NJ tree using JC69 distances (Jukes & Cantor 1969). The rate heterogeneity among sites was assumed to follow a gamma distribution (shape parameter α was ML-estimated) with four categories, each represented by its mean (Yang 1996). The ML-estimated parameters and the model are given in the legend of Fig. 1.

The tree topology found with ML of the pooled mitochondrial data (16S & *coxI*) was not significantly different from the topology inferred from the nuclear PK gene if assessed by the two-tailed K-H-test ($-\ln L_{(\text{treePK}|\text{PKdata})} = 618.49$; $-\ln L_{(\text{tree16S\&coxI}|\text{PKdata})} = 632.50$; $P =$

0.16) with re-estimation of maximum likelihood by non-parametric bootstrapping (RELL) (Kishino & Hasegawa 1989; Kishino et al. 1990). Thus, we pooled the nuclear and mitochondrial data, leaving 1,095 bp (475 bp of the *coxI* and 318 bp of the PK, excluding gaps and ambiguities).

In a further step, the molecular clock hypothesis (i.e., equal rates across all sequences) was tested using the χ^2 approximated likelihood ratio test statistics with OTU's-2 degrees of freedom ($df = 22-2 = 20$) which was rejected with a P -value < 0.01 (Huelsenbeck & Crandall 1997). We explored the tree space by 100 heuristic tree searches using the Tree-Bisection-Reconnection (TBR) algorithm and by randomizing the order of the sequence input in PAUP*. Phylogenetic trees were rooted using *Androctonus australis* as an outgroup. Statistical confidence of phylogenies was assessed using the bootstrap procedure (1,000 pseudoreplicates) (Felsenstein 1985) using PAUP*. These distances were usually $\sim 15\%$ between *Mesobuthus* species and go up to $\sim 20\%$ if *Mesobuthus* sequences were compared to the outgroup.

RESULTS AND DISCUSSION

The three analyzed DNA fragments contained considerable polymorphism among but also within species as estimated using Nei & Li's (1979) nucleotide diversity π and the amount of synonymous versus non-synonymous substitutions of coding regions (Jukes & Cantor 1967) (Table 2). We found 153 mutations out of 475 bp (of which 29 were replace-

ment changes) in the *cox1* fragment, and 19 mutations out of 318 bp (of which 8 were replacement changes) in the PK gene (excluding the outgroup). The 16S fragment contained 131 polymorphic sites out of 302 bp.

The recovered phylogeny (Fig. 1) showed high support for all four included species. This phylogeny allows to address several important taxonomic and evolutionary issues pertaining to the genus *Mesobuthus*. First of all, the phylogeny demonstrates a deep split between the “western clade” of *M. gibbosus* and *M. cyprius* and the “eastern clade” of *M. eupeus* and *M. caucasicus* (Fig. 1); each clade is well supported (100% and 92% bootstrap, respectively). Within the “eastern clade,” there is a strong support for currently accepted species *M. eupeus* (87%) and *M. caucasicus* (94%). The *Mesobuthus caucasicus* sequences are nested within the genus *Mesobuthus* as a sister group to *M. eupeus*. This observation is important as the generic placement of *M. caucasicus* has been controversial since Farzanpay (1987) created a separate monotypic genus *Olivierus* for this species. As our molecular data show, this new genus is paraphyletic with respect to *Mesobuthus* (Fig. 1). Fet & Lowe (2000) listed *Olivierus* as a valid genus but noticed that it was created without any solid justification or revision. The only characters (number of granules on movable finger of pedipalp chela) that Farzanpay (1987) briefly quoted (in Farsi) in support of *Olivierus*, were borrowed from Vachon (1958) species-level descriptions, and are not diagnostic for a genus. Further, our molecular data do not support monophyly of *Olivierus*. Therefore, we propose to list this genus as a synonym of *Mesobuthus*, as was traditionally accepted by all authors before Farzanpay (1987): *Olivierus* Farzanpay 1987 = *Mesobuthus* Vachon 1950, NEW SYNONYMY.

Another observation refers to the subspecific structure of *M. eupeus*, the type species of the genus *Mesobuthus*. This species is extremely polymorphic, and has 14 formally valid subspecies ranging from Turkey to China (Vachon 1958; Fet 1989; Fet & Lowe 2000), with most subspecies described from Iran. The nominotypic subspecies *M. e. eupeus* (C.L. Koch 1839) is found in the Caucasus and Turkey, while most populations from Central Asia are classified as *M. e. thersites* (C.L. Koch, 1839). In our analysis, the nominotypic

subspecies (Turkey) is highly supported (bootstrap 100%) while the Central Asian sequences present a more complicated case. A separate clade exists for a sand desert population from Turkmenistan (MeuTUa1), as opposed to the clade of several populations from Uzbekistan, Kazakhstan and China (bootstrap 68%). This can be an indication of an ancient separation between southern and northern desert forms, possibly valid at least at the subspecies level (Fet 1994). The further phylogenetic analysis could result in elevating these subspecies to the species level. It is interesting that already Birula (1917) grouped all subspecies of *M. eupeus* into two species groups (“sections”), “*eupeus*” and “*thersites*”; however, status of these groups was never examined.

Within *M. caucasicus*, a very strongly supported clade (bootstrap 98%) groups populations from central Uzbekistan (Bukhara, Fergana) and Kazakhstan, while those from southern Uzbekistan (Jarkurgan, Babatag) group outside. It remains to be seen if genetic separation in this case is matched by the morphological variation, as there are several subspecies described from Central Asia as well.

In addition, in our phylogeny the northern Central Asian populations of both *Mesobuthus eupeus* and *M. caucasicus* (Kazakhstan) appear to be derived compared to the southern populations of both species (Uzbekistan); this could be the result of progressive Tertiary aridization and spreading of the arid scorpion species from south to north to the sand and clay deserts (Fet 1994).

The presented data also allow a calibration of a molecular clock using the separation of Cyprus from the Anatolian mainland (5.2 Mya) after the Messinian salinity crisis, during which gene flow between island and mainland populations could have been possible. The Mediterranean Basin was refilled within only 100 yrs, which provides an excellent calibration point for a molecular clock (Hsü et al. 1977; Gantenbein & Largiadèr 2002). Thus, the sequence divergence between Anatolia and Cyprus was estimated to 0.09 ± 0.01 (0.10 ± 0.01 for 16S), which results in a sequence divergence rate of 0.017 per My. This rate estimate is somewhat higher than previous estimates in scorpions for *Mesobuthus gibbosus* (Gantenbein & Largiadèr 2002) but lies absolutely in the range of scorpions such

as *Buthus occitanus* and *Centruroides* (Buthidae) and other invertebrates such as butterflies, beetles and crickets (Brower 1994; Fleischer et al. 1998; Gómez-Zurita et al. 2000; Gantenbein et al. 2001; Gantenbein & Lariadèr 2003).

The genus *Mesobuthus* was created when Vachon (1950) initiated a large-scale “splitting” revision of the traditional genus *Buthus* Leach 1815. Its composition is still controversial. For instance, several Indian species are classified currently in this *Mesobuthus* but their generic identity is unclear (Fet & Lowe 2000). Separate species have been only recently described from Cyprus (Gantenbein et al. 2000b) and confirmed for northern Israel (Fet et al. 2000). Numerous subspecies exist in *M. eupeus* and *M. caucasicus* but morphological characters are inconclusive as for their relationship and taxonomic status. Further application of new molecular markers will facilitate our understanding of taxonomy and evolution of this common scorpion genus.

ACKNOWLEDGMENTS

We are thankful to our late colleagues, Gary A. Polis (California, USA) and Chingiz Tarabaev (Almaty, Kazakhstan) for their help and support in the scorpion work. We dedicate this contribution to their lives, which inspired many young biologists all over the world. We are grateful to A. Scholl and A. Davidson for donation of the specimens used in this study, and to Mark D. Barker and W. Ian Towler for their skilled help in the lab. B.G. was supported by a SNF grant (SNF-83EU-065528) and the Royal Society (R-36579) and V.F. by a National Geographic Society Research and Exploration Fund (grant 7001-0001). V.F. and A.V.G. are grateful to Alex and Elena Kreuzberg, Viktor Lukarevsky, Alexander Tarabrin, Gochmurat Kutlyev and Sergei Morozov for their hospitality and help in field logistics.

LITERATURE CITED

- Birula, A. (Byalynitskii-Birulya, A.A.). 1917. Arachnoidea Arthrogastra Caucasia. Pars I. Scorpiones. Zap. Kavkazskogo Muz. (Mém. Mus. Caucase), Imprimerie de la Chancellerie du Comité pour la Transcaucasie, Tiflis, A(5), 253 pp. (in Russian); English translation by Israel Program for Scientific Translations, Jerusalem 1964, 170 pp.
- Brower, A.V.Z. 1994. Rapid morphological radiation and convergence among races of the butterfly *Heliconius erato* inferred from patterns of mitochondrial DNA evolution. Proceedings of the National Academy of Sciences USA. 91:6491–6495.
- Farzanpay, R. 1987. (Knowing scorpions). No. 312, Biology 4, Central University Publications, Teheran, 231 pp. (in Farsi, with Latin index).
- Felsenstein, J. 1985. Confidence limits on phylogenies: An approach using the bootstrap. Evolution 39:783–791.
- Fet, V. 1989. A catalogue of scorpions (Chelicerata: Scorpiones) of the USSR. Rivista di Museo Civico Scienze Naturali E. Caffi (Bergamo) 13:73–171.
- Fet, V. 1994. Fauna and zoogeography of scorpions (Arachnida: Scorpiones) in Turkmenistan. Pp. 525–534. In Biogeography and Ecology of Turkmenistan (Monographiae Biologicae 72) (V. Fet & K.I. Atamuradov, eds.). Kluwer Academic Publishers, Dordrecht-Boston.
- Fet, V., B. Gantenbein, A.V. Gromov, G. Lowe & W.R. Lourenço. 2003. The first molecular phylogeny of Buthidae (Scorpiones). Euscorpius 4: 1–12.
- Fet, V., B.E. Hendrixson, W.D. Sissom & G. Levy. 2000. First record for the genus *Mesobuthus* Vachon, 1950 in Israel: *Mesobuthus nigrocinctus* (Ehrenberg, 1828), comb. n. (Scorpiones: Buthidae) from Mt. Hermon. Israel Journal of Zoology 46:157–169.
- Fet, V. & G. Lowe. 2000. Family Buthidae C. L. Koch, 1837. Pp. 54–286. In Catalog of the Scorpions of the World (1758–1998) (V. Fet, W.D. Sissom, G. Lowe & M.E. Braunwalder). New York Entomological Society, New York.
- Fet, V., G.A. Polis & W.D. Sissom. 1998. Life in sandy deserts: the scorpion model. Journal of Arid Environments 39:609–622.
- Fet, V., M.E. Soleglad & M.D. Barker. 2001. Phylogenetic analysis of the “*hirsutus*” group of the genus *Hadrurus* Thorell (Scorpiones: Iuridae) based on morphology and mitochondrial DNA. Pp. 139–161. In Scorpions 2001. In Memoriam Gary A. Polis (V. Fet & P.A. Selden, eds.). British Arachnological Society, Burnham Beeches, Bucks, UK.
- Fleischer, R.C., C.E. McIntosh & C.L. Tarr. 1998. Evolution on a volcanic conveyor belt: using phylogeographic reconstructions and K-Ar-based ages of the Hawaiian Islands to estimate molecular evolutionary rates. Molecular Ecology 7: 533–545.
- Folmer, O., M. Black, W. Hoeh, R. Lutz & R. Vrijenhoek. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. Molecular Marine Biology and Biotechnology 3:294–299.
- Gantenbein, B., V. Fet, M. Barker & A. Scholl.

- 2000a. Nuclear and mitochondrial markers suggest the existence of two parapatric scorpion species in the Alps: *Euscorpius germanus* (C. L. Koch, 1837) and *E. alpha* di Caporiacco, 1950, stat. nov. (Scorpiones, Euscorpiidae). *Revue Suisse de Zoologie* 107:843–869.
- Gantenbein, B., V. Fet & M.D. Barker. 2001. Mitochondrial DNA markers reveal a deep, divergent phylogeny in *Centruroides exilicauda* (Wood, 1863) (Scorpiones: Buthidae). Pp. 235–253. *In* *Scorpions 2001*. In *Memoriam Gary A. Polis* (V. Fet & P.A. Selden, eds.). British Arachnological Society, Burnham Beeches, Bucks, UK.
- Gantenbein, B., V. Fet, C.R. Largiadèr & A. Scholl. 1999a. First DNA phylogeny of the genus *Euscorpius* Thorell, 1876 (Scorpiones, Euscorpiidae) and its bearing on the taxonomy and biogeography of this genus. *Biographica* (Paris) 75:59–72.
- Gantenbein, B., C. Kropf, C.R. Largiadèr, & A. Scholl. 2000b. Molecular and morphological evidence for the presence of a new buthid taxon (Scorpiones: Buthidae) on the Island of Cyprus. *Revue Suisse de Zoologie* 107:213–232.
- Gantenbein, B., C.R. Largiadèr & A. Scholl. 1999b. Nuclear and mitochondrial gene variation of *Buthus occitanus* (Amoreux, 1789) across the Strait of Gibraltar. *Revue Suisse de Zoologie* 106:760.
- Gantenbein, B. & C.R. Largiadèr. 2002. *Mesobuthus gibbosus* (Scorpiones: Buthidae) on the island of Rhodes—Hybridisation between Ulysses' stowaways and native scorpions? *Molecular Ecology* 11:925–938.
- Gantenbein, B. & C.R. Largiadèr. 2003. The phylogeographic importance of the Strait of Gibraltar as a gene flow barrier in terrestrial arthropods: A case study with the scorpion *Buthus occitanus* as model organism. *Molecular Phylogenetics and Evolution*. 28:119–130.
- Gómez-Zurita, J., C. Juan & E. Petitpierre. 2000. The evolutionary history of the genus *Timarcha* (Coleoptera, Chrysomelidae) inferred from mitochondrial COII gene and partial 16S rDNA sequences. *Molecular Phylogenetics & Evolution* 14:304–317.
- Gromov, A.V. 2001. On the northern boundary of scorpions (Arachnida: Scorpiones) in Central Asia. Pp. 301–307. *In* *Scorpions 2001*. In *Memoriam Gary A. Polis* (V. Fet & P.A. Selden, eds.). British Arachnological Society, Burnham Beeches, Bucks, UK.
- Huelsenbeck, J.P. & K.A. Crandall. 1997. Phylogeny estimation and hypothesis testing using maximum likelihood. *Annual Review of Ecology and Systematics* 28:437–66.
- Huelsenbeck, J.P. & B. Rannala. 1997. Phylogenetic methods come of age: testing hypotheses in an evolutionary context. *Science* 276:227–232.
- Hsü, K.J., L. Montadert, D. Bernoulli, M.B. Cita, A. Erickson, R.E. Garrison, R.B. Kidd, F. Mèlières, C. Müller & R. Wright. 1977. History of the Mediterranean salinity crisis. *Nature* 267:399–403.
- Jukes, T.H. & C.R. Cantor. 1969. Evolution of protein molecules. Pp. 21–132. *In* *Mammalian Protein Metabolism* (Munro, H. N., ed.). Academic Press, New York.
- Kishino, H. & M. Hasegawa. 1989. Evaluation of the maximum likelihood estimate of the evolutionary tree topologies from DNA sequence data, and the branching order in Hominoidea. *Journal of Molecular Evolution* 29:170–179.
- Kishino, H., T. Miyata & M. Hasegawa. 1990. Maximum likelihood inference of protein phylogeny and the origin of chloroplasts. *Journal of Molecular Evolution* 30:151–160.
- Nei, M. & W. H. Li. 1979. Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proceedings of the National Academic Sciences of the USA* 76:5269–5273.
- Posada, D. & K.A. Crandall. 1998. MODELTEST: testing the model of DNA substitution. *Bioinformatics* 14:817–818.
- Sambrook, J., E.F. Fritsch & T. Maniatis. 1989. *Molecular Cloning: a Laboratory Manual*. Cold Harbor Laboratory Press, New York.
- Simon, C., F. Frati, A. Beckenbach, B. Crespi, H. Liu & P. Flook. 1994. Evolution, weighting, and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain reaction primers. *Annals of the Entomological Society of America* 87:651–701.
- Swofford, D.L. 1998. PAUP* Phylogenetic analysis using parsimony (*and other methods). Version 4. Sinauer Associates, Sunderland, MA.
- Swofford, D.L., G.J. Olsen, P.J. Waddell & D.M. Hillis. 1996. Phylogenetic inference. Pp. 407–510. *In* *Molecular systematics* (Hillis, D.M., Moritz, C. & Mable, B.K., eds.). Sinauer Associates, Sunderland, MA.
- Tamura, K. & M. Nei. 1993. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology & Evolution* 10:512–526.
- Thompson, J.D., T.J. Gibson, F. Plewniak, F. Jeanmougin & D.G. Higgins. 1997. The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research* 24:4876–4882.
- Towler, W.I., J. Ponce Saavedra, B. Gantenbein & V. Fet. 2001. Mitochondrial DNA reveals divergent phylogeny in tropical *Centruroides* (Scorpiones: Buthidae) from Mexico. *Biographica* (Paris) 77:157–172.
- Vachon, M. 1950. Études sur les Scorpions. III (suite). *Déscription des Scorpions du Nord de*

- l'Afrique. Archives de l'Institut Pasteur d'Algérie 28:152–216.
- Vachon, M. 1952. Etudes sur les scorpions. Institut Pasteur d'Algérie, Alger, 482 pp.
- Vachon, M. 1958. Scorpionidea (Chelicerata) de l'Afghanistan. The 3rd Danish Expedition to Central Asia. (Zoological Results 23). Videnskabelige Meddelelser fra Dansk Naturhistorisk Forening i København 120:121–187.
- Yang, Z. 1996. Among-site rate variation and its impact on phylogenetic analyses. Trends in Ecology & Evolution 11:367–372.
- Manuscript received 1 June 2001, revised 7 April 2003.*