

Methods to Identify Silk Gland Activation Patterns in Spider Spinning Behaviors

Maitry Jani¹, Martín Ramírez² and Jonas O. Wolff^{1,3}: ¹Evolutionary Biomechanics, Zoological Institute and Museum, University of Greifswald, Loitzer Str. 26, 17489 Greifswald, Germany; E-mail: maitry.jani@uni-greifswald.de; ²Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”, Consejo Nacional de Investigaciones Científicas y Técnicas, Buenos Aires, Argentina; ³School of Natural Sciences, Macquarie University, Sydney, NSW 2109, Australia

Abstract. Spiders possess multiple types of silk glands, producing silk materials with contrasting properties, which are deployed in distinct behavioral contexts, such as locomotion, prey capture and egg casing. Whereas the diversity of silk glands and spigots across different spider families is relatively well described, their biological functions (i.e., with which behavior each gland type is activated) are poorly known. Here we provide an overview about available methods and approaches to determine the biological function of spider silk glands, and evaluate their advantages and disadvantages based on our experiences and a test study on *Pholcus phalangioides* (Fuesslin, 1775) (Pholcidae) and *Kukulcania hibernalis* (Hentz, 1842) (Filistatidae).

Keywords: Spider silk, silk spinning-behavior, silk types, multi-method approach
<https://doi.org/10.1636/JoA-S-24-016>

Unlike other silk-producing animals, spiders deploy silk in multiple ways throughout their life. From predation to reproduction, dispersal, and maternal care, spiders use silk as a multifunctional tool (Apstein 1889; Gorb & Barth 1994; Blackledge et al. 2005; Gheysens et al. 2005; Foelix 2011; Bell et al. 2005; Blackledge 2012). With the potential for up to eight distinct types of silk glands (excluding the epiandrous glands found in adult males), each presumably serving one or more specific biological functions and specialized to produce fibers and glues with different mechanical properties (Blackledge & Hayashi 2006; Foelix 2011), spiders possess a sophisticated apparatus to spin a variety of fiber-based materials. The silk glands are present in the opisthosoma and open through specialized tube- or nozzle-like cuticular protrusions, the spigots, that are located on modified abdominal appendages, the spinnerets (Foelix 2011; Eberhard 2020). The morphological arrangement of spigots on the spinnerets, combined with varied utilization of different silk glands across different behaviors, is distinct between different spider families and has captivated arachnologists for years. Spiders’ spinning organs are very diverse yet sufficiently stereotyped, so that spider systematics has benefitted from their morphology to infer evolutionary relationship (e.g., Coddington 1989; Platnick et al. 1991; Griswold et al. 2005).

Since the beginning of spider studies, a large variety of silk materials and spinning behaviors has been described, along with the morphological variation of gland types, numbers and spigot arrangements (Apstein 1889; Coddington 1989; Eberhard 2010; Alfaro et al. 2018). However, in only very few cases was it observed which glands are involved in which behavioral contexts. This is a significant problem for the general understanding of how the morphology of the spinning apparatus, silk material composition and silk property variation have co-evolved with the behavioral ecology in spiders.

A reason for this knowledge gap is a lack of standardized protocols, resulting in a patchwork of multiple methods that are not well documented and have not been evaluated regarding their efficiency. Some studies have relied on studying silk samples collected from spider webs or by intercepting specific spinning behaviors. These samples were then analyzed with polarized light microscopy (PLM) or electron microscopy to distinguish fibers based on diameters or optical properties (Work 1981; Blackledge et al. 2005; Wolff et al. 2017).

Some approaches, including the paraffin embedding method pioneered by Peters in 1982, have been repeatedly used in behavioral studies (Peters 1990, 1992; Gorb & Barth 1996; Townley & Tillinghast 2003; Grannemann et al. 2019). Other, less explored approaches to identify the use of specific silk glands in specific spider behaviors, include snap freezing the spider followed by cryo-scanning electron microscopy, high speed video recording, or the experimental blocking of specific spigots (Barth et al. 1998; Wolff 2019; Wolff et al. 2015, 2024). Another promising approach is the comparison of the chemical profiles of silk products and gland contents (Lefèvre et al. 2011; Wolff et al. 2024).

Thus far, there is no study that combined these different techniques in the same system. The present study presents an overview of the available set of methodologies to identify gland use in different behavioral contexts or the glandular origin of silk products, with the aim to give a practical guide for experimentalists.

As a test object, we studied the behaviorally versatile cosmopolitan cellar spider (*Pholcus phalangioides* (Fuesslin, 1775)). This species possesses two anterior lateral spinnerets each with one major ampullate gland spigot, and two posterior median spinnerets each with one minor ampullate and one aciniform gland spigot. These glands produce fibers of different widths. In addition, there are one large piriform and multiple small piriform gland spigots on each anterior lateral spinneret, producing glue-like silk secretions (Apstein 1889; Kirchner 1986; Kooor 1986). The posterior lateral spinnerets do not bear any functional spigots in this species (Platnick et al. 1991). In addition to tests with *P. phalangioides*, paraffin experiments and silk reeling tests were performed on Southern house spiders (*Kukulcania hibernalis* (Hentz, 1842)), which are cribellate spiders possessing a more complex spinning apparatus with seven different types of silk glands.

METHODS AND RESULTS

Direct observation assisted by high-magnification lenses and high-speed camera.—Ideally it can be directly observed from which spigot(s) a given silk product is coming. However, this is possible to observe only in a limited number of spinning behaviors due to the extremely small size of spigots, thin diameter and translucence of silk fibers (making them hard to see), and




Figure 1.—Single frames of high-speed videos of spinning *Pholcus phalangioides*. Arrowheads highlight silk fiber origination. (A) Filming of dragline initiation with high magnification lens, permitting focus on the spinning apparatus (details on method in Wolff 2021). Here it is visible, that both minor ampullate and aciniform fibers originated from posterior median spinnerets and could be included in the dragline. (B) Posterior close-up view on the spider walking in an inverted position and leaving behind a dragline (arrowhead), where fibers were drawn from all three glands (major ampullate, minor ampullate and aciniform glands), as indicated by the fibers' number, different thickness and origination from both anterior and posterior median spinnerets (detail enhanced in inset). (C) Prey immobilization wrapping behavior, where fibers are drawn with the help of 4th pair of legs; note that two strands of silk can be distinguished. *Abbreviations:* Ac – aciniform gland silk fiber; ALS – anterior lateral spinneret; MA – major ampullate gland silk fiber; miA – minor ampullate gland silk fiber; PMS – posterior median spinneret.

often high speed of movements during silk spinning. Direct observation can best be done in behaviors that only depend on the choreography of the spinnerets and can be performed while the spider is immobilized. One of us (JOW) has previously described a method that makes it possible to directly observe the spigots that are active during the production of silk anchor points and dragline initiation by letting a spider spin against a glass slide within a confined, highly magnified camera field of view (Wolff 2019, 2021; Fig. 1A). High-speed footage was also used to understand the spigot origin and kinematics of spinning the ribbon-like silk of recluse spiders of the sicariid genus *Loxosceles* Heineken & Lowe, 1832 (Magalhaes et al. 2017), and the silk gland usage in the wrap-attacks of Gnaphosidae (Wolff et al. 2017).

Methodology: Here we tested how well a high-speed camera (Phantom MIRO LC 320S) equipped with a macro-lens (Canon 100mm) may resolve fiber origination in free moving spiders. For this, videos were recorded while the spiders exhibited behaviors such as dragline spinning, prey wrapping, and bridging. Recordings were done at 2000 frames per second. The camera was placed on a tripod with rollers for enhanced mobility to easily follow the spider and keep it in focus. Recordings were then replayed at low frame rates (slow motion) to discern from which spinnerets fibers were coming (Fig. 1 B & C).

Evaluation: The advantage of this method is that silk emergence can be directly observed. It also allows us to infer information on the spinning kinematics, for example at what speed fibers are spun.

The disadvantage of this method is that it requires expensive equipment and strong illumination, which might inhibit natural behaviors in many nocturnal species. Further, only in behaviors where most of the spider body stays stationary (e.g., cribellar combing; Grannemann et al. 2019) may it be possible to focus on the spinning apparatus with a lens that allows sufficient magnification to distinguish between spigots. In those cases where the spider's body remains stationary, we found that it may be feasible to observe from which spinnerets fibers are coming and, in some cases, how many fibers are included into the silk product. In the case of *P. phalangioides*, due to the low number of spigots and

comparably large differences between fiber diameters, it was possible to indirectly infer which silk gland was used (i.e., observing from which spinneret silk was coming and then associating diameters of fibers in the final silk products with spigot sizes). However, the necessary magnification to resolve these details came at the cost of a low depth of field, making it impossible to continuously keep the opisthosoma in focus during rapid spinning, such as prey wrapping behavior. Notably, rare behaviors, such as the construction of gumfoot lines or egg sacs, which always happen at night, could not be captured with this method.

Paraffin-fixation.—Due to the microscopic scale, to enhance clarity in silk fiber emergence from spigots, ideally a scanning electron microscope (SEM) is used. However, as animals (usually) cannot be observed *in vivo* in the SEM, a useful approach is to fix the spinning behavior. One way to achieve this is to pour hot paraffin onto the spinning apparatus and the spider. The paraffin solidifies and embeds the spinnerets and fibers, fixing their position. The spider is then transferred to ethanol or a chemical fixative, the paraffin chemically dissolved, and the sample dehydrated, sputter coated and mounted for SEM. This technique was successfully used to infer the origin of airborne ballooning and bridging lines (Peters 1990), draglines (Gorb & Barth 1996; Townley & Tillinghast 2003), and the fibers composing the cribellar bands (Peters 1982, 1984, 1992).

Sample preparation: One of us (MJR) has applied the paraffin method to the Southern house spider *Kukulcania hibernalis* (see Grannemann et al. 2019). The melted paraffin was stored in a 50 ml Falcon vial immersed in an insulated container (thermos) with hot water, then transferred to a smaller vial to drop on the spinning spiders. The spiders were transferred to 95% ethanol for a week, the paraffin was then dissolved with xylol, and the spider was then put through a graded series to 100% ethanol, critical point dried and sputter coated with Au/Pd, and finally observed using standard (FEI-XL30TMP) or field emission electron scanning microscope (Zeiss Supra). The rate of success was moderate and disparate for different fiber types, ca. 1 successful fixation per 3–5 trials for cribellar or minor ampullate fibers, but 1 out of 10 for the paracribellar and major ampullate fibers.




Figure 2.—SEM images from paraffin and cryo fixation methods. Arrowheads highlight silk fiber origination. (A–C) Paraffin fixation of *Kukulcania hibernalis*. (A) Fixation of anchor spinning behavior, showing fibers emerging from piriform and major ampullate glands. (B) Fixation of cribellar combing behavior, showing fibers originating from cribellum spigots and (C) fibers from paracribellar spigots. (D–F) Cryo fixation images of *Pholcus phalangioides* while producing (D) dragline, where major ampullate glands were active, (E) bridging line, where minor ampullate glands were active, and (F) dragline anchorage, where both the large piriform gland (emitting glue to attach fibers to the substrate) and major ampullate glands were active. *Abbreviations:* Cr – cribellar spigot; MA – major ampullate spigot; MiA – minor ampullate spigot; PC – paracribellar spigot; Pi – piriform spigot.

Results: The technique allowed us to image short stretches of fibers coming out of the spigots (Fig. 2 A–C). The subsequent matching of those fibers to the natural web structures was done indirectly by comparing fiber morphology and diameter. The matching of ampullate and cribellar fibers was clear, because of their disparate diameters and characteristic morphology: the cribellar fibers were ultra-thin, major ampullate fibers were thick, and minor ampullate fibers were undulating, very thick and oval in cross-section. Paracribellar fibers, on the other hand, were tentatively matched by diameter alone. Finally, in addition to

individual fiber morphology and diameter, the number of spigots observed in SEM preparations of the spinning organs corresponded with the fibers found in the cribellate band and its supporting thread: two minor ampullate gland spigots corresponded to the thick undulating fibers, six paracribellar gland spigots with the thinner undulating fibers, many cribellar gland spigots with the nano-fibrils, and six major ampullate gland spigots with the fibers of the supporting thread.

Evaluation: The advantage of this method is that fiber origin can be directly observed. The technique can be applied in the

field by transporting the molten paraffin in an insulated container. The disadvantages are the time-consuming sample preparation and high error rate, i.e., it requires the killing of many spiders to obtain suitable results. It is therefore not suitable for species that occur in low abundance nor for rarely observed behaviors.

Cryo-fixation.—The idea of this approach is the same as with the paraffin method: the instant fixation of a given behavior to enable the microscopic observation of which spigots emitted which silks. Liquid nitrogen is used for fixation, freezing the spider in an instant. These frozen specimens can then be transferred directly to a Cryo-scanning electron microscope (Wolff et al. 2015) or fixed in chilled ethanol and investigated with classical SEM as described below.

Experimental design. Here, two different experimental setups were employed to study the dragline formation. For draglines produced while walking, the individuals were allowed to walk on a traverse steel stand wrapped with multiple layers of tissue paper. As the spider walked, it laid down a dragline behind it. The tissue paper, along with attached dragline and spider, was carefully lifted with wooden forceps and submerged in a liquid nitrogen container. The cryo-fixed spider was then retrieved from the liquid nitrogen using the same wooden forceps and preserved in chilled ethanol (see “Sample preservation, preparation and investigation” for further steps).

The dragline spun during abseiling behavior was studied by letting the spider descend from an elevated surface (rod or vial) and directly plunged it in a liquid nitrogen container. Dispersal behavior, characterized by the use of an airborne bridging line (Huber 2023), was investigated by confining individuals to a small stand, coating the surrounding area with Vaseline to restrict movement, and submerging the stand in liquid nitrogen when spiders started to show bridging behavior.

To study the prey-wrapping behavior, each individual was housed in 15 × 10 × 10 cm plastic boxes for 1 to 2 weeks during which a web was constructed. The housing container was opened on the front and placed in a larger polystyrene box to prevent spillage during liquid nitrogen application. Then one blowfly was presented to the spider with forceps, which was then attacked and wrapped. Once prey wrapping was initiated, liquid nitrogen was immediately poured onto the spider to snap-freeze it.

Sample preservation, preparation and investigation.—Cryo-fixed spiders should be immediately preserved in chilled ethanol, or there is a chance that the abdomen bursts and leaks, rendering any further investigation impossible.

In our trials, the cryo-fixed spiders were mounted into ethanol that was stored at -80°C . The specimens were then stored at -80°C or -40°C (no difference in the outcome was observed between the two storage temperatures). Prior to sample preparation, the ethanol concentration was gradually increased to 96%. The specimens were then transferred to freezers with increasing temperatures (-40°C , -10°C), and finally to the lab at room temperature. The spiders were then subjected to critical point drying and affixed to the aluminium stubs with conductive silver glue. With scanning electron microscopy (SEM), spigots were examined for the presence of silk fibers coming from the spigot pores, indicative of gland activity during the observed behaviors.

Evaluation. The advantage of this approach is that it permits the direct visualization of silk gland activity during behaviors. We could observe not only the emergence of fibers (Fig. 2 D–F), but also the addition of sticky silk from the large piriform glands

in silk anchor spinning (Fig. 2F). The sample preparation for this method is less time-consuming than for the paraffin method. If direct transfer to a Cryo-SEM is performed, it should be even possible to observe the emergence of viscid silk products, such as aggregate silk, that are soluble in polar liquids and would get lost during chemical fixation procedures.

The disadvantages are that it is a destructive method (the spider dies and can only be observed once). Usually, a high number of spiders is needed as the silk emergence is visible only in a subset of the samples. The technique requires lab facilities to store and handle liquid nitrogen. Therefore, in contrast to the paraffin technique, it cannot be used in the field. In comparison to other approaches, the cryo-fixation method followed by SEM is comparably time-consuming and labor-intensive. Furthermore, we found that the freezing sometimes leads to squeezing out of excess liquid silk in spigots, forming a tiny blob or fiber piece coming out of spigots. While regularly spun fibers usually show a smooth and even shape, sometimes it is difficult to interpret if material seen coming from the spigot was an artifact due to the method.

Experimental spigot blocking.—This follows the classical approach in functional morphology to identify the role of a certain structure in a certain behavioral context, by modifying or removing the structure and investigating how the behavioral performance changes. By blocking (sealing) or destroying specified spigots, the production of this silk by the spider is inhibited. By comparing with control spiders with unblocked spigots or observations made on the same spiders before the experimental manipulation, it is then determined, which silk materials change their composition or which silk-based behaviors are altered or not displayed any longer.

Anaesthesia and preparation: Live individuals of *P. phalangioides* were anesthetized with CO_2 . Subsequently, the spiders were immobilized with the ventral side upwards on a soft Styrofoam platform using masking tape, allowing for clear observation of the spinnerets under a stereomicroscope. To ensure that the spiders are not completely unable to spin silk structures, care was taken not to block the large piriform and major ampullate spigots, which are required for dragline production and anchorage and are the prerequisite for construction of silk scaffolds. Instead, we tried to target the spigots of the posterior median spinnerets or the small piriform spigots.

Spigot blocking procedure: Various materials, including nail polish, eyelash glue, Cyanoacrylate super glue, and water-based (Elmer’s) glue were tested to block specific spigots. The suitability of each material in targeting individual spigots, durability of adhesion and likelihood of contaminating other spigots due to fluid properties was assessed.

Nail polish: The advantage is the bright colour making it easy to clearly see even small amounts, but due to its comparably high viscosity, nail polish proved challenging to be applied in small amounts to target specific spigots. Small amounts tended to not stick very well on the cuticle, and could only be secured by further spreading the nail polish along the lateral edge of the spinneret. On the other hand, the viscosity prevented the contamination of other spigots. In many cases, the spiders were able to remove the nail polish using their hind legs within the two days after treatment. This could be mitigated by applying larger amounts of nail polish, which, however, may cause uneven weight distribution, affecting spider mobility.




Figure 3.—SEM (A–D) and PLM (E–G) images for spigot pore size measurements and fiber diameter measurements of *Pholcus phalangioides*. (A) Major ampullate spigot pore. (B) Minor ampullate spigot pore. (C) Dragline showing two fibers. (D) Detail of gumfoot line (tangle capture line from web), showing all three silk types, distinguishable by fiber width. (E–G) Fibers of different widths observed in egg sac and prey wrap samples. Abbreviations: Ac – aciniform silk fiber; MA – major ampullate silk fiber; MiA – minor ampullate silk fiber.

Eyelash glue: The eyelash glue was sufficiently sticky, but exhibited a slow drying time. This caused problems when the anaesthesia wore off and spiders started to move their spinnerets, leading to cross-contamination.

Super glue: Due to its initially low viscosity, even tiny amounts of super glue were spreading across all spinnerets, causing the accidental blockage of multiple spigots.

Water-based glue: This adhesive had a comparably high viscosity and extended drying period, and was therefore ineffective for applying it to specific spigots.

Evaluation: It was found that it was barely possible to block specific spigots or spigot fields. If small enough amounts were used, the spiders were able to remove the block. The spider's ability to remove adhesive blocks may be taxon-specific. *Pholcus phalangioides* is known to exhibit a thick coating of fluid hydrocarbon on their cuticle, which might act as a repellent to adhesives (Frutiger & Kropf 2019). If higher amounts of glue were used, it contaminated other spigots. Overall, we found that spiders were highly stressed by the experimental treatment and their behavior was greatly affected, even if only single spigots were blocked. Treated spiders usually stopped spinning webs and draglines altogether, making it impossible to determine changes in gland-specific behaviors. This caveat might be species-specific, and it remains unclear if the experimental approach may be effective for other spider taxa.

Correlating fiber and spigot pore diameters.—This approach is based on the observation that the spigots of different gland types often differ in the diameter of their terminal pore opening. Assuming that the spigot cuticle is relatively stiff, small pores would not be able to emit thick fibers, and hence fibers of thicker diameters originate from spigots with larger diameters. As the spigot pores are in the micron to sub-micron scale their investigation using the SEM is required. Fibers may either be visualised with an SEM or polarised light microscopy (PLM) (Blackledge et al. 2005). When comparing the diameters, it is important to keep in mind that silks, and potentially cuticle, too, might slightly shrink in the SEM (Greco et al. 2023).

Methodology: In our case study, seven different silk products (drop-down dragline, walking dragline, bridging line, gumfoot line, web tangle, prey wrap, and egg sac) were collected from multiple individuals on sample holders as described by Ramírez et al. (2013). The opisthosomata of the same individuals were prepared for SEM (sample preparation was performed as mentioned above, except for the temperature gradient steps).

High-resolution images of the spigots were captured, and subsequently each spigot pore diameter was measured using the ImageJ software (Schneider et al. 2012). Spigot pore dimensions were correlated with fiber diameters to establish relationships between spigots and their respective silk products.

Results: In *P. phalangioides*, we found that each spigot type emits a distinct fiber type. The fibers from different silk samples were visually distinguished as thin, thick, and medium sized using the PLM (Fig. 3 E–G). In the SEM, the aciniform spigot pore showed as the smallest, the major ampullate as medium sized (Fig. 3A) and the minor ampullate as the largest (Fig. 3B). However, we found that the minor ampullate spigot had a flexible cuticle surrounding the pore, and in the SEM samples the pore opening was often collapsed, making it hard to quantify the effective diameter (in these cases the average of the long and the short diameter across the oval or slit-like pore was taken). By comparison with the results from the Cryo-method, we found that this spigot ejects fibers with a range of diameters, some of which may be indistinguishable from the diameters of major ampullate and aciniform silk fibers. These characteristics may be more specific to Pholcidae, and less of a problem for spiders of other families.

Evaluation. The advantage of the approach to correlate fibers with spigots based on diameters is that with PLM a large number of silk samples can be analysed in a comparably short time.

The limitation is that matching fibers and spigots based on diameters must be regarded as a hypothesis. It cannot be fully excluded that large pores can emit thin fibers, as not only the duct and pore dimensions, but also the spinning (drawing) speed of fibers from the spigots during particular behaviors may affect the fiber diameter size (Vollrath & Edmonds 1989; Kaplan et al. 1994). Furthermore, the approach is only applicable if spigots of different silk types are clearly distinct in their diameters and exhibit a stable pore shape. Especially in families where more than four silk gland types are present, it might be difficult to distinguish fiber types based on diameters alone, and additional characteristics might have to be taken into account (e.g., birefringence, Raman spectra, etc., see below).

Distinguishing gland products by chemical analysis.—Different silk gland products differ in their chemical composition and base protein (spidroin) structure. By comparing signatures in silk chemistry between the contents of the identified glands and that of the silk product it is possible to determine, which gland was involved in the production of which material. There is a large range of techniques available, such as mass spectroscopy,

Nuclear Magnetic Resonance spectroscopy or X-ray diffraction (Blamires et al. 2023; Wolff et al. 2023).

Sequencing cDNAs generated from captured mRNA isolated from specific glands and the transcript of predicted protein sequences can allow to identify spidroins associated with fiber types (Correa-Garhwal et al. 2019). For instance, this approach was used in the identification of Pyriform Spidroin 1 (PySp1) as the main silk protein used in anchoring dragline silk fibers in the attachment discs of *Latrodectus hesperus* Chamberlin & Ivie, 1935 (Blasingame et al. 2009). As many spidroins are expressed in their highest concentration in specific silk glands, information on spidroin sequences can be used to predict the mass spectra of digested silk fibers. The mass spectra from digested silk fibers of unknown gland origin can then be matched with the reference database (Casem et al. 1999; Hu et al. 2006, 2007; Chaw et al. 2015).

Another approach is the determination of the ratio of different amino acids in the silk sample, as different silk types are known to differ in amino acid composition, and different silk gland types differ in their histochemical characteristics. Andersen (1970) provided some of the earliest application of this techniques and Casem et al. (1999) used amino acid analysis to demonstrate that the content of tubuliform glands of *Araneus diadematus* Clerck, 1757 and *Latrodectus hesperus* are much more serine-rich than their other glands, correlating with the serine-rich egg case fibers, and the serine-rich amino acid sequence of Tubuliform Spidroin (TuSp) (Garb & Hayashi 2005).

However, most of the aforementioned biochemical techniques usually require comparably high amounts of material. Further, these approaches are only applicable if the analysed chemical compounds are highly contrasting in different gland product (as the case for the distinguishing high serine content in tubuliform silk).

Raman spectroscopy is especially promising as it allows the recording of chemical profiles directly in the native sample with high spatial resolution. By obtaining Raman profiles on both the silk products and the content of dissected glands (silk dope), it is, in principle, possible to identify the glandular origin of the silk product. However, it is important to note that the Raman profiles differ between pre- and post-spinning silks due to structural changes occur during the spinning process (Lefèvre et al. 2011). The efficiency of this approach highly depends on how much the Raman profiles of the gland products differ. We have previously used the technique to determine, which of diverse adhesive silks produced by *Pholcus phalangioides* likely are the same gland products (Wolff et al. 2024). In this case, the approach proved very useful in distinguishing the silks. However, we could not obtain the corresponding profiles of gland contents as it was not possible to clearly separate the silk dope from the surrounding epithelia, which probably is only feasible for large species.

Distinguishing gland products by mechanical properties.— Different gland products are known to differ in their mechanical properties (Blackledge & Hayashi 2006). Especially ultimate strain (indicator of how much a material can be stretched before it breaks) and Young's modulus (an indicator of material stiffness) differ between major ampullate, minor ampullate, aciniform and flagelliform silks (Blackledge & Hayashi 2006). These characteristics may be used to distinguish whether sampled silk fibers are made from the same material or not. While the estimation of silk strength and Young's modulus requires a sophisticated setup

(such as a universal testing machine with high resolution load cells), silk strain can be relatively easily quantified by attaching the silk sample to the forks of callipers and stretching out the silk until it breaks.

However, there are strong limitations to the approach of determining the glandular origin of silk materials based on mechanical properties. First, silk properties are almost exclusively known for major ampullate silk, and knowledge on the tensile properties of other silks is extremely limited. In order to create a reference database, it is necessary to do tensile test of isolated silk fibers whose glandular origin is known. This can be done by forcibly silking spiders, where in a restrained spider, specific spigots are brushed with a fine probe that may lead to silk release that is slowly pulled into a strand that is then transferred to the sample holder (Work & Emerson 1982). Our experience with this technique is mixed. In *P. phalangioides*, distinct silk samples can be obtained that way, but is often difficult to identify from which spigot the silk is coming, even if using a stereo microscope with high magnification. We were able to forcibly pull single aciniform and major ampullate fibers, but the minor ampullate fibers tended to stick to aciniform fibers and it was not possible to obtain isolated minor ampullate silk samples with this method. In *Kukulkania hibernalis* we found silking was almost impossible, as the spinning field was inaccessible due to the specific spinneret morphology and setal field covering the spigots in rest.

Second, tensile properties are affected by the way they are spun (Greco et al. 2023) and it is likely that behaviors, in which spiders rapidly pull fibers from the glands (such as during prey wrapping) result in different mechanical properties than behaviors in which spiders slowly and steadily extend the silk (such as in web building), even if the used silk material is the same. Spinning effects on tensile properties seem to be more expressed in some spider taxa than in others, and were previously found insignificant for ampullate silks of *Pholcus phalangioides* (Boutry et al. 2011). However, we have found that bridging lines, that are pulled by *P. phalangioides* at high speed, have different and more variable tensile properties than draglines, even if used materials are the same (Jani et al. 2025). And third, spiders often combine multiple silks in their threads and sheets, and such material mixture creates complex mechanical behavior, in which the properties of the single components may be masked (Liprandi et al. 2024).

Finally, tensile testing is a destructive technique, and therefore cannot be used on rare and valuable samples.

DISCUSSION: COMPARATIVE EVALUATION OF THE METHODOLOGY

We found that each method offered their own advantages, challenges and limitations (summary in Table 1). None of the methods are applicable for all behaviors, silk products or species, therefore we propose a combination of methods as the best approach for a more comprehensive understanding of spider silk production and gland functionality. While SEM and PLM remain essential tools for the detailed structural analysis of silk products and spigots, our observation suggests that relying solely on fiber diameter measurements may not provide a complete understanding on which silk types (gland products) are included in a given silk material. Flexible spigot openings and variation in fiber production rates can lead to variation in fiber diameters (Madsen

Table 1.—Overview on approaches to identify spider silk gland use in behavioral contexts.

Method	Advantages	Disadvantages
Behavioral observation aided by high-speed camera/high-magnification lenses	<ul style="list-style-type: none"> • Direct observation of silk emergence. • May also provide insight into spinning kinematics, including speed of fibre spinning. • Can infer information on silk gland usage and fibre inclusion. 	<ul style="list-style-type: none"> • Requires expensive equipment and strong illumination. • May inhibit natural behaviors, especially in nocturnal species. • Applicability to only a limited set of behaviors. • Low depth of field may make continuous focus challenging during rapid spinning.
Paraffin fixation	<ul style="list-style-type: none"> • Direct observation of fibre origin. • Field applicability with molten paraffin in portable insulated container. 	<ul style="list-style-type: none"> • Time-consuming sample preparation. • High error rate and moderate success, requiring multiple trials. • Requires killing of spiders, unsuitable for low abundance species or rare behaviors.
Cryo-fixation	<ul style="list-style-type: none"> • Direct visualization of silk gland activity during behaviors. • Less time-consuming sample preparation compared to paraffin method. • Permits observation of both solid and viscid silks. 	<ul style="list-style-type: none"> • Destructive method, spiders die and can only be observed once. • Requires a high number of spiders due to high error rate. • Requires lab facilities for liquid nitrogen handling. • Not suitable for field use. • Time-consuming and labor-intensive compared to other methods. • Freezing may lead to squeezing out of excess liquid silk, potentially creating artifacts.
Spigot blocking	<ul style="list-style-type: none"> • Experimental approach to infer the role of specific glands in different behaviors, including such that cannot be directly observed. 	<ul style="list-style-type: none"> • High stress on spiders during experimental treatment. • Limited success in blocking specific spigots or spigot fields. • Spiders may remove adhesive blocks, especially smaller amounts (potential taxon-specific differences in ability to remove blocks). • Contamination of other spigots if higher amounts of adhesive used. • Applicability may vary among spider species.
Correlating fibre and spigot pore diameters	<ul style="list-style-type: none"> • Allows correlation of spigots to fibres even when not functional at the moment of fixation. • Allows the inference of gland origin for silk products, whose construction is difficult to observe (such as egg cases or nocturnal web construction). • Large number of silk samples can be analysed comparatively quickly with polarized light microscopy (PLM) 	<ul style="list-style-type: none"> • Matching of fibres and spigots based on diameters is a hypothesis and may not always be accurate. • Possibility that large pores emit thin fibres may challenge correlation. • Applicability limited to cases where spigots of different silk types exhibit clearly distinct diameters and stable pore shapes. • Difficult for species with many silk gland types. • Additional characteristics such as birefringence or Raman spectra may be necessary for distinguishing silk types.
Distinguishing silk types by chemical analysis	<ul style="list-style-type: none"> • Allows the inference of gland origin for silk products, whose construction is difficult to observe. • May also inform about differences in silk composition and/or protein structure. • Raman spectroscopy provides high spatial resolution chemical profiles directly in native samples. • Can be applied to both solid and viscid silks. 	<ul style="list-style-type: none"> • Most techniques (except Raman spectroscopy) require high amounts of material, which may be unfeasible to collect. • Raman profiles differ between pre- and post-spinning silks due to structural changes, which may challenge correlations. • Efficiency depends on the degree of difference in Raman profiles of different gland products. • Difficulty in obtaining profiles of gland contents due to challenges in separating silk dope from surrounding epithelia, particularly in smaller species.
Distinguishing silk types by mechanical properties	<ul style="list-style-type: none"> • Allows the inference of gland origin for silk products, whose construction is difficult to observe. • Also informs about the mechanical properties of silk lines, with ecological implications. • Silk strain can be relatively quantified without expensive equipment. 	<ul style="list-style-type: none"> • Limited knowledge on tensile properties of non-major ampullate silks. • Requires a reference database with silks whose glandular origin is known (e.g., collection by forcibly silking, which can be challenging in some species and/or gland types). • Tensile properties may be affected by spinning speed and behaviors (such effects may vary between silk types and taxa). • Spiders often combine multiple silks, masking the properties of individual gland products. • Destructive technique, unsuitable for rare or valuable samples.

et al. 1999; Vollrath & Köhler 1996). This may lead to discrepancies and false assumptions about thin fiber origins (Vollrath & Edmonds 1989). Therefore, it is recommended that gland uses are further studied via direct observation methods, like high-speed videography and cryo-fixation, where possible. Even if silk emergence cannot be observed directly, often the movement of the animals or their body parts can be informative to logically infer fiber origins in conjunction with the number and diameters of the fibers in the spun silk material, and the number, sizes and arrangement of spigots (Eberhard 2010).

As we progressively learn from direct observations such as silking, video recording, paraffin- and cryo-fixation, it becomes possible to extrapolate such knowledge to related species, so that fiber origin can be in certain cases safely inferred from indirect observations, such as fiber diameter and morphology, and their disposition in compound structures such as cribellar bands, viscid threads and silk anchors.

In conclusion, our study underscores the importance of adopting a multi-method approach to study spider behavior and silk gland utilization effectively. By combining direct observation methods with the microscopy of silk products, researchers can gain deeper insights into the complex dynamics of spider silk production, behavioral ecology and gland functionality.

ACKNOWLEDGMENTS

We thank Paula Heinz for her help with animal care and sample collection. We thank Gabriele Uhl for access to light microscopes and lab facilities. We thank Rabea Schlüter from the Imaging Center of the Biology unit at the University of Greifswald for her support with SEM microscopy. This work was funded by an ERC Starting Grant to JOW (101040724 — SuPer-Silk) under the European Union's Horizon 2020 research and innovation programme.

LITERATURE CITED

- Alfaro RE, Griswold CE, Miller KB. 2018. Comparative spigot ontogeny across the spider tree of life. *PeerJ* 6:e4233.
- Andersen SO. 1970. Amino acid composition of spider silks. *Comparative Biochemistry and Physiology* 35:705–711.
- Apstein C. 1889. Bau und Funktion der Spinnrüsen der Araneida. *Archive der Naturgeschichte* 55:29–74.
- Barth FG, Gorb SN, Landolfi MA. 1998. Dragline-associated behavior of the orb web spider *Nephila clavipes* (Araneioidea, Tetragnathidae). *Journal of Zoology* 244:323–330.
- Bell JR, Bohan DA, Shaw EM, Weyman GS. 2005. Ballooning dispersal using silk: world fauna, phylogenies, genetics and models. *Bulletin of Entomological Research* 95:69–114.
- Blackledge TA. 2012. Spider silk: a brief review and prospectus on research linking biomechanics and ecology in draglines and orb webs. *Journal of Arachnology* 40:1–12.
- Blackledge TA, Hayashi CY. 2006. Silken toolkits: biomechanics of silk fibers spun by the orb web spider *Argiope argentata* (Fabricius 1775). *Journal of Experimental Biology* 209:2452–2461.
- Blackledge TA, Cardullo RA, Hayashi CY. 2005. Polarized light microscopy, variability in spider silk diameters, and the mechanical characterization of spider silk. *Invertebrate Biology* 124:165–173.
- Blamires SJ, Rawal A, Edwards AD, Yarger JL, Oberst S, Allardycy BJ, et al. 2023. Methods for silk property analyses across structural hierarchies and scales. *Molecules* 28:2120.
- Blasingame E, Tuton-Blasingame T, Larkin L, Falick AM, Zhao L, Fong J, et al. 2009. Pyriform spidroin 1, a novel member of the silk gene family that anchors dragline silk fibers in attachment discs of the black widow spider, *Latrodectus hesperus*. *Journal of Biological Chemistry* 284:29097–29108.
- Boutry C, Řezáč M, Blackledge TA. 2011. Plasticity in major ampullate silk production in relation to spider phylogeny and ecology. *PLoS One* 6:e22467.
- Casem ML, Turner D, Houchin K. 1999. Protein and amino acid composition of silks from the cob weaver, *Latrodectus hesperus* (black widow). *International Journal of Biological Macromolecules* 24:103–108.
- Chaw RC, Correa-Garhwal SM, Clarke TH, Ayoub NA, Hayashi CY. 2015. Proteomic evidence for components of spider silk synthesis from black widow silk glands and fibers. *Journal of Proteome Research* 14:4223–4231.
- Coddington JA. 1989. Spinneret silk spigot morphology: evidence for the monophyly of orbweaving spiders, Cyrtophorinae (Araneidae), and the group Theridiidae plus Nesticidae. *Journal of Arachnology* 17:71–95.
- Correa-Garhwal SM, Clarke TH, Janssen M, Crevecoeur L, McQuillan BN, Simpson AH, et al. 2019. Spidroins and silk fibers of aquatic spiders. *Scientific Reports* 9:13656.
- Eberhard WG. 2010. Possible functional significance of spigot placement on the spinnerets of spiders. *Journal of Arachnology* 38:407–414.
- Eberhard WG. 2020. Spider Webs: Behavior, Function, and Evolution. University of Chicago Press. Chicago, London.
- Foelix R. 2011. Biology of Spiders. 3rd edition. Oxford University Press, New York.
- Frutiger M, Kropf C. 2019. An anti-adhesive surface coating reduces adhesion during contact with cribellar threads in *Pholcus phalangioides* (Araneae, Pholcidae) but not in the web-owning spider *Uloborus plumipes* (Araneae, Uloboridae). *BioRxiv*:838250.
- Garb JE, Hayashi CY. 2005. Modular evolution of egg case silk genes across orb-weaving spider superfamilies. *Proceedings of the National Academy of Sciences* 102:11379–11384.
- Gheysens T, Beladjal L, Gellynck K, Van Nimmen E, Van Langenhove L, Mertens J. 2005. Egg sac structure of *Zygiella x-notata* (Arachnida, Araneidae). *Journal of Arachnology* 33:549–57.
- Gorb SN, Barth FG. 1996. A new mechanosensory organ on the anterior spinnerets of the spider *Cupiennius salei* (Araneae, Ctenidae). *Zoology* 116:7–14.
- Gorb SN, Barth FG. 1994. Locomotor behavior during prey-capture of a fishing spider, *Dolomedes plantarius* (Araneae: Araneidae): galloping and stopping. *Journal of Arachnology* 22:89–93.
- Grannemann CC, Meyer M, Reinhardt M, Ramírez MJ, Herberstein ME, Joel AC. 2019. Small behavioral adaptations enable more effective prey capture by producing 3D-structured spider threads. *Scientific Reports* 9:17273.
- Greco G, Schmuck B, Jalali SK, Pugno NM, Rising A. 2023. Influence of experimental methods on the mechanical properties of silk fibers: A systematic literature review and future road map. *Biophysics Reviews* 4:031301.
- Griswold CE, Ramírez MJ, Coddington J, Platnick N. 2005. Atlas of phylogenetic data for entelegyne spiders (Araneae: Araneomorphae: Entelegynae) with comments on their phylogeny. *Proceedings of the California Academy of Sciences* 4:1–324.
- Hu X, Kohler K, Falick AM, Moore AM, Jones PR, Vierra C. 2006. Spider egg case core fibers: trimeric complexes assembled from TuSp1, ECP-1, and ECP-2. *Biochemistry* 45:3506–3516.
- Hu X, Yuan J, Wang X, Vasanthavada K, Falick AM, Jones PR, et al. 2007. Analysis of aqueous glue coating proteins on the silk fibers of the cob weaver, *Latrodectus hesperus*. *Biochemistry* 46:3294–3303.
- Huber BA. 2023. Do pholcid spiders balloon? *Arachnology* 19:885–887.
- Jani M, Promnil S, Heinz P, Liprandi D, Wolff JO. 2025. Functional diversity and behavioural use of fibre producing silk glands in the

- cosmopolitan cellar spider *Pholcus phalangioides* (Fuesslin, 1775). *Journal of Experimental Biology*, in press.
- Kaplan D, Adams WW, Farmer B, Viney C. 1994. *Silk: Biology, Structure, Properties, and Genetics*. ACS Publications, Washington DC.
- Kirchner VW. 1986. Das Netz der Zitterspinne (*Pholcus phalangioides* Fuesslin)(Araneae: Pholcidae). *Zoologischer Anzeiger* 216:151–169.
- Kovoor J. 1986. Affinités de quelques Pholcidae (Araneae) décelables d'après les caractères de l'appareil séricigène. *Mémoires de la Société Entomologique de Belgique* 33:111–118.
- Lefèvre T, Boudreault S, Cloutier C, Pézolet M. 2011. Diversity of molecular transformations involved in the formation of spider silks. *Journal of Molecular Biology* 405:238–253.
- Liprandi D, Ramírez M, Schlüter S, Baumgart L, Joel AC, Michalik P, Wolff JO. 2024. Hierarchical looping results in extreme extensibility of silk fiber composites produced by Southern house spiders (*Kukulcania hibernalis*). *Interface Focus* 14:20230071
- Madsen B, Shao ZZ, Vollrath F. 1999. Variability in the mechanical properties of spider silks on three levels: interspecific, intraspecific and intraindividual. *International Journal of Biological Macromolecules* 24:301–306.
- Magalhaes IL, Ravelo AM, Scioscia CL, Peretti AV, Michalik P, Ramírez MJ. 2017. Recluse spiders produce flattened silk rapidly using a highly modified, self-sufficient spinning apparatus. *Journal of Zoology* 303: 27–35.
- Peters HM. 1982. Wie Spinnen der Familie Uloboridae ihre Beute einspinnen und verzehren. *Verhandlungen des Naturwissenschaftlichen Vereins in Hamburg (NF)* 25:142167.
- Peters HM. 1984. The spinning apparatus of Uloboridae in relation to the structure and construction of capture threads (Arachnida, Araneida). *Zoomorphology* 104:96–104.
- Peters HM. 1990. On the structure and glandular origin of bridging lines used by spiders for moving to distant places. *Acta Zoologica Fennica* 190:309–314.
- Peters HM. 1992. Über Struktur und Herstellung von Fangfaden cribellater Spinnen der Familie Eresidae (Arachnida, Araneae). *Verhandlungen des Naturwissenschaftlichen Vereins in Hamburg* 33:213–227.
- Platnick NI, Coddington JA, Forster RR, Griswold CE. 1991. Spinneret morphology and the phylogeny of haplogyne spiders (Araneae, Araneomorphae). *American Museum Novitates* 3016:1–73.
- Ramírez MJ, Ravelo AM, Lopardo L. 2013. A simple device to collect, store and study samples of two-dimensional spider webs. *Zootaxa* 3750:189–192.
- Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nature Methods* 9:671–675.
- Townley MA, Tillinghast EK. 2003. On the use of ampullate gland silks by wolf spiders (Araneae, Lycosidae) for attaching the egg sac to the spinnerets and a proposal for defining nubbins and tartipores. *Journal of Arachnology* 31:209–245.
- Vollrath F, Edmonds DT. 1989. Modulation of the mechanical properties of spider silk by coating with water. *Nature* 340:305–307.
- Vollrath F, Köhler T. 1996. Mechanics of silk produced by loaded spiders. *Proceedings of the Royal Society of London. Series B: Biological Sciences* 263:387–391.
- Wolff JO. 2019. The evolution of dragline initiation in spiders: multiple transitions from multi- to single-gland usage. *Diversity* 12:4.
- Wolff JO. 2021. Evolutionary kinematics of spinneret movements for rapid silk thread anchorage in spiders. *Journal of Comparative Physiology A* 207:141–152.
- Wolff JO, Ashley LJ, Schmitt C, Heu C, Denkova D, Jani M, et al. 2024. From fibers to adhesives: evolution of spider capture threads from web anchors by radical changes in silk gland function. *Journal of the Royal Society Interface* 21:20240123.
- Wolff JO, Cherry BR, Yarger JL, Adler L, Thomas DS, Hook JM, et al. 2023. Organic salt composition of pressure sensitive adhesives produced by spiders. *Frontiers in Ecology and Evolution* 11:1123614.
- Wolff JO, Grawe I, Wirth M, Karstedt A, Gorb SN. 2015. Spider's super-glue: thread anchors are composite adhesives with synergistic hierarchical organization. *Soft Matter* 11:2394–2403.
- Wolff JO, Řezáč M, Krejčí T, Gorb SN. 2017. Hunting with sticky tape: functional shift in silk glands of araneophagous ground spiders (Gnaphosidae). *Journal of Experimental Biology* 220:2250–2259.
- Work RW. 1981. Web components associated with the major ampullate silk fibers of orb-web-building spiders. *Transactions of the American Microscopical Society* 100:1–20.
- Work RW, Emerson PD. 1982. An apparatus and technique for the forcible silking of spiders. *Journal of Arachnology* 10:1–10.

Manuscript received 30 May 2024, revised 21 October 2024, accepted 23 October 2024.